

Case/Application number: 10/551,292  
Priority Filing Date: 4/15/2003  
Format for Search Results: Short  
Matching of unusual acronyms or initials:

Identify the novelty:

See limitation in amended claim 1 regarding X and 54. "b molecule" may be searched open-ended.


Additional comments:

Please search the structure in claim 1 amended 4/20/2003. See also specific structure in claim 25.

10/551,292

=> d que stat l7

L5 STR



VAR G1=14/10/12

VAR G2=15/16/19/21

VAR G3=14/10/12

REP G4=(1-2) C

REP G5=(1-2) C

VAR G6=S/N/P

NODE ATTRIBUTES:

CONNECT IS E2 RC AT 14

CONNECT IS E2 RC AT 15

DEFAULT MLEVEL IS ATOM

GGCAT IS UNS AT 13

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 26

STEREO ATTRIBUTES: NONE

L7 827 SEA FILE=REGISTRY SSS FUL L5


100.0% PROCESSED 767106 ITERATIONS

827 ANSWERS

SEARCH TIME: 00.01.01

=> d que stat l12

L5 STR



VAR G1=14/10/12

VAR G2=15/16/19/21


VAR G3=14/10/12

10/551,292

REP G4=(1-2) C  
 REP G5=(1-2) C  
 VAR G6=S/N/P  
 NODE ATTRIBUTES:  
 CONNECT IS E2 RC AT 14  
 CONNECT IS E2 RC AT 15  
 DEFAULT MLEVEL IS ATOM  
 GGCAT IS UNS AT 13  
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:  
 RING(S) ARE ISOLATED OR EMBEDDED  
 NUMBER OF NODES IS 26

STEREO ATTRIBUTES: NONE  
 L7 827 SEA FILE=REGISTRY SSS FUL L5  
 L10 STR



VAR G1=14/10/12  
 VAR G2=15/16/19/21  
 VAR G3=14/10/12  
 REP G4=(1-2) CH2  
 REP G5=(1-2) CH2  
 VAR G6=S/N/P  
 NODE ATTRIBUTES:  
 CONNECT IS E1 RC AT 11  
 CONNECT IS E2 RC AT 14  
 CONNECT IS E2 RC AT 15  
 CONNECT IS E2 RC AT 22  
 DEFAULT MLEVEL IS ATOM  
 GGCAT IS UNS AT 13  
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:  
 RING(S) ARE ISOLATED OR EMBEDDED  
 NUMBER OF NODES IS 26

STEREO ATTRIBUTES: NONE  
 L12 90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10

100.0% PROCESSED 827 ITERATIONS  
 SEARCH TIME: 00.00.01

90 ANSWERS

=> d que nos 131  
 L5 STR

```


L7      827 SEA FILE=REGISTRY SSS FUL L5
L10     STR
L12     90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10
L14     QUE SPE=ON ABB=ON PLU=ON SANTOS, I?/AU,AUTH
L15     QUE SPE=ON ABB=ON PLU=ON REGO, I?/AU,AUTH
L16     QUE SPE=ON ABB=ON PLU=ON CORREIA, J?/AU,AUTH
L17     QUE SPE=ON ABB=ON PLU=ON PAULO, A?/AU,AUTH
L18     QUE SPE=ON ABB=ON PLU=ON ALVES, S?/AU,AUTH
L19     QUE SPE=ON ABB=ON PLU=ON VITOR, R?/AU,AUTH
L20     QUE SPE=ON ABB=ON PLU=ON MALLINCKRODT/CS, SO, PA
L21     QUE SPE=ON ABB=ON PLU=ON AY<2004 OR PY<2004 OR PRY<20
      04 OR MY<2004 OR REVIEW/DT
L22     QUE SPE=ON ABB=ON PLU=ON BIOMOLECUL? OR (BIO(1W)MOLEC
      UL?) OR (BIOLOGIC?(3A)MOLECUL?)
L23     QUE SPE=ON ABB=ON PLU=ON CHELAT?
L24     QUE SPE=ON ABB=ON PLU=ON "CHELATING AGENTS"+PFT,OLD,N
      EW,NT/CT
L25     46 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L12
L26     19 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L25 AND (L22 OR L23
      OR L24)
L27     46 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON (L25 OR L26)
L28     18 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L27 AND (L14 OR L15
      OR L16 OR L17 OR L18 OR L19 OR L20)
L29     28 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L27 NOT L28
L30     23 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L29 AND L21
L31     28 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON (L29 OR L30)

```

```

=> d que stat l33
L10     STR

```



```

VAR G1=14/10/12
VAR G2=15/16/19/21
VAR G3=14/10/12
REP G4=(1-2) CH2
REP G5=(1-2) CH2
VAR G6=S/N/P
NODE ATTRIBUTES:
CONNECT IS E1 RC AT 11
CONNECT IS E2 RC AT 14
CONNECT IS E2 RC AT 15
CONNECT IS E2 RC AT 22
DEFAULT MLEVEL IS ATOM
GGCAT IS UNS AT 13
DEFAULT ECLEVEL IS LIMITED

```

```

GRAPH ATTRIBUTES:

```

10/551,292

RING(S) ARE ISOLATED OR EMBEDDED  
NUMBER OF NODES IS 26

STEREO ATTRIBUTES: NONE

L33 10 SEA FILE=WPIX SSS FUL L10

100.0% PROCESSED 11053 ITERATIONS

10 ANSWERS

SEARCH TIME: 00.00.13

=> d que nos 138

L10 STR  
L14 QUE SPE=ON ABB=ON PLU=ON SANTOS, I?/AU,AUTH  
L15 QUE SPE=ON ABB=ON PLU=ON REGO, I?/AU,AUTH  
L16 QUE SPE=ON ABB=ON PLU=ON CORREIA, J?/AU,AUTH  
L17 QUE SPE=ON ABB=ON PLU=ON PAULO, A?/AU,AUTH  
L18 QUE SPE=ON ABB=ON PLU=ON ALVES, S?/AU,AUTH  
L19 QUE SPE=ON ABB=ON PLU=ON VITOR, R?/AU,AUTH  
L20 QUE SPE=ON ABB=ON PLU=ON MALLINCKRODT/CS,SO,PA  
L22 QUE SPE=ON ABB=ON PLU=ON BIOMOLECUL? OR (BIO(1W)MOLEC  
UL?) OR (BIOLOGIC?(3A)MOLECUL?)  
L23 QUE SPE=ON ABB=ON PLU=ON CHELAT?  
L33 10 SEA FILE=WPIX SSS FUL L10  
L34 3 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON (RABNX7/DCN OR RAFVJB/DCN  
OR RAFVJC/DCN OR RAFVJD/DCN OR RAFVJE/DCN OR RAFVJF/DCN OR  
RAFVJG/DCN OR RAFVJ8/DCN OR RAFVJ9/DCN OR RAMT8E/DCN) OR  
L33/DCR  
L35 2 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON L34 AND (L22 OR L23)  
L36 3 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON (L34 OR L35)  
L37 1 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON L36 AND (L14 OR L15 OR  
L16 OR L17 OR L18 OR L19 OR L20)  
L38 2 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON L36 NOT L37

=> d que nos 139

L5 STR  
L7 827 SEA FILE=REGISTRY SSS FUL L5  
L10 STR  
L12 90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10  
L39 0 SEA FILE=MEDLINE SPE=ON ABB=ON PLU=ON L12

=> d que nos 141

L5 STR  
L7 827 SEA FILE=REGISTRY SSS FUL L5  
L10 STR  
L12 90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10  
L41 0 SEA FILE=EMBASE SPE=ON ABB=ON PLU=ON L12

=> d his 142

(FILE 'BIOSIS, CABA, AGRICOLA, BIOTECHNO, DRUGU, VETU' ENTERED AT  
13:30:22 ON 25 JUN 2009)

L42 0 S L12

=> d que nos 142

L5 STR  
L7 827 SEA FILE=REGISTRY SSS FUL L5

10/551,292

L10 STR  
L12 90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10  
L42 0 SEA L12

=> dup rem 131 138 139 141 142

L39 HAS NO ANSWERS

L41 HAS NO ANSWERS

L42 HAS NO ANSWERS

FILE 'HCAPLUS' ENTERED AT 13:37:35 ON 25 JUN 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIX' ENTERED AT 13:37:35 ON 25 JUN 2009

COPYRIGHT (C) 2009 THOMSON REUTERS

PROCESSING COMPLETED FOR L31

PROCESSING COMPLETED FOR L38

PROCESSING COMPLETED FOR L39

PROCESSING COMPLETED FOR L41

PROCESSING COMPLETED FOR L42

L45 28 DUP REM L31 L38 L39 L41 L42 (2 DUPLICATES REMOVED)

ANSWERS '1-27' FROM FILE HCAPLUS

ANSWER '28' FROM FILE WPIX

=> file stnguide

FILE 'STNGUIDE' ENTERED AT 13:37:48 ON 25 JUN 2009

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Jun 19, 2009 (20090619/UP).

=> fil hcap wpix

FILE 'HCAPLUS' ENTERED AT 13:38:10 ON 25 JUN 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIX' ENTERED AT 13:38:10 ON 25 JUN 2009

COPYRIGHT (C) 2009 THOMSON REUTERS

=> d que 121

L21 QUE SPE=ON ABB=ON PLU=ON AY<2004 OR PY<2004 OR PRY<20  
04 OR MY<2004 OR REVIEW/DT

=> s 145 and 121

'2004' NOT A VALID FIELD CODE

'2004' NOT A VALID FIELD CODE

L46 23 L45 AND L21

=> dup rem 146

PROCESSING COMPLETED FOR L46

L47 23 DUP REM L46 (0 DUPLICATES REMOVED)

ANSWERS '1-22' FROM FILE HCAPLUS

ANSWER '23' FROM FILE WPIX

=> file stnguide

10/551,292

FILE 'STNGUIDE' ENTERED AT 13:39:14 ON 25 JUN 2009  
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Jun 19, 2009 (20090619/UP).

=> d ibib ed abs hitind hitstr 1-22

YOU HAVE REQUESTED DATA FROM FILE 'HCAPLUS, WPIX' - CONTINUE? (Y)/N:y

L47 ANSWER 1 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2000:862644 HCAPLUS Full-text

DOCUMENT NUMBER: 134:246525

TITLE: Syntheses, structure and properties of cobalt-(II) and  
-(III) complexes of pentadentate N4S ligands with  
appended pyrazolyl groups: evidence for  
cobalt(II)-dioxygen reversible binding

AUTHOR(S): Bhattacharyya, Sudeep; Ghosh, Dipesh; Mukhopadhyay,  
Suman; Jensen, William P.; Tiekink, Edward R. T.;  
Chaudhury, Muktimoy

CORPORATE SOURCE: Department of Inorganic Chemistry, Indian Association  
for the Cultivation of Science, Calcutta, 700 032,  
India

SOURCE: Dalton (2000), (24), 4677-4682

CODEN: DALTFG; ISSN: 1470-479X

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 134:246525

ED Entered STN: 11 Dec 2000

AB Cobalt-(II) and -(III) complexes of pentadentate N4S ligands based on Me 2-  
aminocyclopent-1-ene-1-carbodithioate with appended pyrazolyl groups (3,5-  
Me2C3HN2CH2)2NCH(R)CH2NHC5H6C(:S)SCH3 (R = H, Hmmeed; CH3, Hmmpcd) were  
prepared and characterized by IR, 1H NMR and electronic spectroscopy. Two of  
these compds. have also structurally been characterized by x-ray single  
crystal diffraction analyses. Cobalt(II) in [Co(mmpcd)]ClO4 (1) shows a five-  
coordinate, trigonal bipyramidal geometry while its cobalt(III) counterpart,  
[Co(mmpcd)Cl]ClO4 (2), reveals a six-coordinated distorted octahedral  
structure by the inclusion of a chloride ligand in its equatorial plane. In  
DMF or acetonitrile solution, 1 can bind dioxygen reversibly as indicated by  
EPR spectra recorded at cryogenic temps. Metal-dioxygen binding in 1 appears  
to be weak, possibly due to its trigonal bipyramidal structure and the  
presence of a sulfur donor in the ligand framework. Electronic spectra of the  
cobalt(III) complexes show two LMCT bands in the near UV region, tentatively  
assigned to S→CoIII charge transfer.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 75, 77

IT 174280-57-0

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant for preparation of cobalt(III) Me

(bis(dimethylpyrazolylmethyl)aminoethyl)aminocyclopentenecarbodithioato  
complex)


IT 174280-57-0

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant for preparation of cobalt(III) Me

(bis(dimethylpyrazolylmethyl)aminoethyl)aminocyclopentenecarbodithioato  
complex)

RN 174280-57-0 HCAPLUS  
 CN 1-Cyclopentene-1-carbodithioic acid,  
 2-[[2-[bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]amino]ethyl]amino]-,  
 methyl ester (CA INDEX NAME)



REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 2 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2001:43719 HCAPLUS Full-text

DOCUMENT NUMBER: 134:246535

TITLE: Cobalt(II) complexes of ethylenediamine-based pyrazole ligands: the crystal structure of [{N',N'-bis(3,5-dimethylpyrazol-1-ylmethyl)}-N,N-dimethylethylenediamine]cobalt(II) tetraphenylborate  
 AUTHOR(S): Lee, Soon Ae; Lim, Jong Wan; Roh, Soo-Gyun; Yeo, Hwan Jin; Jeong, Jong Hwa

CORPORATE SOURCE: Department of Chemistry, Kyungpook National University, Taegu, 702-701, S. Korea

SOURCE: Bulletin of the Korean Chemical Society (2000), 21(12), 1271-1273

CODEN: BKCSDE; ISSN: 0253-2964

PUBLISHER: Korean Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 134:246535

ED Entered STN: 18 Jan 2001

AB {N',N'-bis(pyrazol-1-ylmethyl)}-N,N-dimethylethylenediamine (L) and {N',N'-bis(3,5-dimethylpyrazol-1-ylmethyl)}-N,N-dimethylethylenediamine (L1) were prepared and complexed with CoCl<sub>2</sub> to give [CoLCl]BPh<sub>4</sub> and [CoL1Cl]BPh<sub>4</sub>. The crystal structure of [CoL1Cl]BPh<sub>4</sub>·0.5Me<sub>2</sub>CO was determined [crystal data: orthorhombic, space group P212121, Z = 8, R1 = 0.0617, wR2 = 0.1488]. The complex has a trigonal bipyramidal structure in which Cl and 1 ethylenediamine N atom are axial.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 28, 75

IT 330153-74-7P 330153-75-8P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and complexation with cobalt)

IT 330153-74-7P 330153-75-8P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and complexation with cobalt)



RN 330153-74-7 HCAPLUS  
 CN 1,2-Ethanediamine, N1,N1-dimethyl-N2,N2-bis(1H-pyrazol-1-ylmethyl)- (CA INDEX NAME)



RN 330153-75-8 HCAPLUS  
 CN 1,2-Ethanediamine, N1,N1-bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-N2,N2-dimethyl- (CA INDEX NAME)



REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 3 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1999:687174 HCAPLUS [Full-text](#)

DOCUMENT NUMBER: 132:44035

TITLE: Palladium(II)-Induced Activation of Carbon-Nitrogen Single Bond of Coordinated N4S Ligand. Characterization of Product with Modified Ligand Structure: Kinetics versus Thermodynamic Considerations

AUTHOR(S): Bhattacharyya, Sudeep; Weakley, Timothy J. R.; Chaudhury, Muktimoy

CORPORATE SOURCE: Department of Inorganic Chemistry, Indian Association for the Cultivation of Science, Calcutta, 700 032, India

SOURCE: Inorganic Chemistry (1999), 38(23), 5453-5456

CODEN: INOCAJ; ISSN: 0020-1669

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 29 Oct 1999

AB Reaction of Pd(OAc)<sub>2</sub> with 2-(β-bis(3,5-dimethylpyrazol-1-ylmethyl)aminoethylamino)cyclopent-1-enedithiocarboxylate (Hmmedc) in MeOH generated two products isolated as perchlorate salts, [Pd(N3S)-CH<sub>2</sub>Me<sub>2</sub>Pz]ClO<sub>4</sub> (1) and [Pd(N3S)-CH<sub>2</sub>OMe]ClO<sub>4</sub> (2). The square-planar mol. structure of 1 was determined by x-ray crystallog. Formation of 2 involves C-N bond cleavage and requires the presence of Pd(II). However, compound 1 once formed does not undergo methanolysis reaction further. Mechanistic implications of the C-N bond cleavage through a structural anomeric effect in an unstable reactive intermediate is discussed.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 67, 75

IT 67-56-1, Methanol, reactions 174280-57-0

RL: RCT (Reactant); RACT (Reactant or reagent)


(palladium(II)-induced activation of carbon-nitrogen single bond of coordinated N4S ligand via methanolysis)

IT 174280-57-0

RL: RCT (Reactant); RACT (Reactant or reagent)

(palladium(II)-induced activation of carbon-nitrogen single bond of coordinated N4S ligand via methanolysis)

RN 174280-57-0 HCAPLUS

CN 1-Cyclopentene-1-carbodithioic acid,  
2-[[2-[bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]amino]ethyl]amino]-,  
methyl ester (CA INDEX NAME)

REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 4 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1999:508196 HCAPLUS Full-text

DOCUMENT NUMBER: 131:214344

TITLE: Heterocyclic substituted silatranes. Part I. Synthesis and characterization of pyrazolyl substituted aminoalkylsilatranes

AUTHOR(S): Nasim, M.; Tharmaraj, P.; Venkataramani, P. S.

CORPORATE SOURCE: Defence Materials and Stores Research and Development Establishment DMSRDE PO, Kanpur, 208013, India

SOURCE: Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry (1999), 29(7), 1249-1263

CODEN: SRIMCN; ISSN: 0094-5714

PUBLISHER: Marcel Dekker, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 16 Aug 1999

AB N-[Bis(3,5-dimethylpyrazol-1-ylmethyl)aminopropyl]- (6a) and N-[bis(3,5-dimethylpyrazol-1-ylmethyl)aminopropyl]-3,7,10- trimethylsilatranes (6b) were obtained in high yields from the reaction of N-[bis(3,5-dimethylpyrazol-1-ylmethyl)aminopropyl]triethoxysilane (3a) with triethanolamine (4a) or triisopropanolamine (4b). The corresponding N-Ph derivative (6c) was also obtained by the reaction of N-phenyl-[(3,5-dimethylpyrazol-1-ylmethyl)aminopropyl]trimethoxysilane (3b) with (4a). N-[3,5-Dimethylpyrazol-1-ylmethyl-3- silatranylaminopropyl]-N'-[bis(3,5-dimethylpyrazol-1-ylmethyl)]ethylenediamine (10a) and N-[3,5-dimethylpyrazol-1-ylmethyl]-3'-[(3,7,10- trimethylsilatranyl)aminopropyl]-N'-[bis(3,5-dimethylpyrazol-1-

ylmethyl)ethylenediamine (10b) also were synthesized by the reaction of N-[3,5-dimethylpyrazol-1-ylmethyl]-N'-[bis(3,5-dimethylpyrazol-1-ylmethyl)ethylenediamine]trimethoxysilane (8) with (4a) and (4b). The precursor alkoxy compds. (3a), (3b) and (8) themselves were synthesized by reaction of 1-hydroxymethyl-3,5-dimethylpyrazole (2) with 3-aminopropyltriethoxy- (1a), N-phenylaminopropyltrimethoxy- (1b) and N-[2-aminoethyl(aminopropyl)]trimethoxysilanes (7), resp.

CC 29-6 (Organometallic and Organometalloidal Compounds)

IT 74789-21-2P 85952-93-8P 137684-04-9P 242464-71-7P 242464-72-8P

242464-74-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(for preparation of pyrazolyl substituted aminoalkylsilatranes)

IT 242464-66-0P 242464-67-1P 242464-68-2P 242464-69-3P

242464-70-6P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of pyrazolyl substituted aminoalkylsilatranes)


IT 242464-74-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(for preparation of pyrazolyl substituted aminoalkylsilatranes)

RN 242464-74-0 HCAPLUS

CN 1,2-Ethanediamine, N1,N1,N2-tris[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-N2-[3-(trimethoxysilyl)propyl]- (CA INDEX NAME)





IT 242464-69-3P 242464-70-6P

RL: SPN (Synthetic preparation); PREP (Preparation)



(preparation of pyrazolyl substituted aminoalkylsilatranes)

RN 242464-69-3 HCAPLUS

CN 1,2-Ethanediamine, N1,N1,N2-tris[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-N2-[3-(2,8,9-trioxa-5-aza-1-silabicyclo[3.3.3]undec-1-yl)propyl]- (CA INDEX NAME)




12




REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 5 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1999:65917 HCAPLUS Full-text  
 DOCUMENT NUMBER: 130:190922  
 TITLE: Nickel(II) in an N4S Donor Environment: An Unprecedented Alcoholysis Reaction through the Activation of a Carbon-Nitrogen Single Bond  
 AUTHOR(S): Bhattacharyya, Sudeep; Weakley, Timothy J. R.; Chaudhury, Muktimoy  
 CORPORATE SOURCE: Department of Inorganic Chemistry, Indian Association for the Cultivation of Science, Calcutta, 700 032, India  
 SOURCE: Inorganic Chemistry (1999), 38(4), 633-638  
 CODEN: INOCAJ; ISSN: 0020-1669  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 ED Entered STN: 02 Feb 1999  
 GI



- AB Pentadentate N4S ligands based on Me 2-aminocyclopent-1-ene-1-dithiocarboxylate with flexible pyrazolyl arms (Me2pzCH2)2NC2H3RNHC5H6CSSSCH3 (R = H, Hmmeed and R = Me, Hmmpcd) undergo a nickel(II)-induced alcoholysis reaction through the activation of a saturated C-N bond linkage. The products obtained are square-planar complexes [(I); R = H, R' = Me (1); R = Me, R' = Me, Et, n-Pr (2-4)] containing a modified ligand structure possessing an N3S donor set and a pendant arm that holds the alkoxy group provided by the solvent. [Ni(N3S)-CH2OMe]ClO4 (1) crystallizes in the triclinic space group P.hivin.1 with a 10.4886(5), b 10.706(1), c 11.487(1) Å, α 108.784(4), β 108.887(6), γ 95.139(6)°, and Z = 2; while [Ni(N3S)'-CH2OPr]ClO4 (4) has the monoclinic space group P21/n with a 8.875(2), b 18.629(2), c 15.399(2) Å, β 91.37(2)°, and Z = 4 per unit cell. Complexes 1-4 with acyclic ligand environments have interesting electrochem. behavior in acetonitrile, involving a reversible Ni(II)/Ni(I) reduction, E1/2 .apprx.-1.0 V, and a Ni(II)/Ni(III) irreversible oxidation, Epa .apprx.1.0 V vs. Ag/AgCl as the reference. The coulometrically reduced solution of 2 displays a rhombic EPR spectrum at 77 K characteristic of nickel(I) with g1 = 2.217, g2 = 2.170, and g3 = 2.054.
- CC 78-7 (Inorganic Chemicals and Reactions)  
Section cross-reference(s): 72, 75
- IT 64-17-5, Ethanol, reactions 67-56-1, Methanol, reactions 71-23-8, Propanol, reactions 174280-57-0 174280-58-1  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(nickel promoted alcoholysis of pyrazolylmethyl C-N bond in preparation of (pyrazolylmethylaminoalkyl)aminocyclopentenedithiocarboxylato complexes)
- IT 174280-57-0  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(nickel promoted alcoholysis of pyrazolylmethyl C-N bond in preparation of (pyrazolylmethylaminoalkyl)aminocyclopentenedithiocarboxylato complexes)
- RN 174280-57-0 HCAPLUS
- CN 1-Cyclopentene-1-carbodithioic acid,  
2-[[2-[bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]amino]ethyl]amino]-,  
methyl ester (CA INDEX NAME)



REFERENCE COUNT: 63 THERE ARE 63 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 6 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2000:237932 HCAPLUS Full-text  
Correction of: 1997:64019

DOCUMENT NUMBER: 132:245490  
Correction of: 126:219991

TITLE: Ion-pair extraction behavior of transition metal(II) cations as charged complexes with ethylenediamine derivatives having heterocyclic pendant arms

AUTHOR(S): Hirayama, Naoki; Iimuro, Shinji; Kubono, Koji; Kokusen, Hisao; Honjo, Takaharu

CORPORATE SOURCE: Kanazawa University, Kanazawa, 920-11, Japan

SOURCE: Analytica Chimica Acta (1997), 339(1-2), 115-121

CODEN: ACACAM; ISSN: 0003-2670

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 13 Apr 2000

AB The use of N,N,N',N'-tetrakis(1'-pyrazolylmethyl)-1,2-diaminoethane (tpzen), N,N,N',N'-tetrakis(3',5'-dimethylpyrazol-1'-ylmethyl)-1,2-diaminoethane (Me8tpzen) and N,N,N',N'-tetrakis(2'-pyridylmethyl)-1,2-diaminoethane (tpen) as complexation reagents for ion-pair extraction of metal(II) cations into nitrobenzene was studied. The order of extractability of the metals was tpen >> Me8tpzen > tpen. Although these mols. have four nitrogen-containing heterocyclic pendant arms and act as sexadentate ligands normally in aqueous solution, the result of numerical anal. concerning the extraction behavior indicated that they act as bidentate ligands in the extraction system. The pendant arms were not concerned in chelate formation and the arms acted to raise the hydrophobicity and the extractability of the complexes.

CC 79-4 (Inorganic Analytical Chemistry)

Section cross-reference(s): 68

IT 85264-42-2P 85264-43-3P

RL: ARG (Analytical reagent use); PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); ANST (Analytical study); PREP (Preparation); PROC (Process); USES (Uses)

(preparation and dissociation constant and use in ion-pair extraction of transition metal(II) cations)

IT 85264-42-2P 85264-43-3P


RL: ARG (Analytical reagent use); PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); ANST (Analytical study); PREP (Preparation); PROC (Process); USES (Uses)

(preparation and dissociation constant and use in ion-pair extraction of transition

metal(II) cations)


RN 85264-42-2 HCAPLUS

CN 1,2-Ethanediamine, N1,N1,N2,N2-tetrakis(1H-pyrazol-1-ylmethyl)- (CA INDEX NAME)



RN 85264-43-3 HCAPLUS

CN 1,2-Ethanediamine, N1,N1,N2,N2-tetrakis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]- (CA INDEX NAME)



L47 ANSWER 7 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1996:145239 HCAPLUS [Full-text](#)

DOCUMENT NUMBER: 124:218506

ORIGINAL REFERENCE NO.: 124:40069a,40072a


TITLE: Zinc(II) and Copper(II) Complexes of Pentacoordinating (N4S) ligands with Flexible Pyrazolyl Arms: Syntheses, Structure, Redox, and Spectroscopic Properties

AUTHOR(S): Bhattacharyya, Sudeep; Kumar, Sujit Baran; Dutta, Subodh Kanti; Tiekink, Edward R. T.; Chaudhury, Muktimoy

CORPORATE SOURCE: Department of Inorganic Chemistry, Indian Association for the Cultivation of Science, Calcutta, 700 032,



India  
 SOURCE: Inorganic Chemistry (1996), 35(7), 1967-73  
 CODEN: INOCAJ; ISSN: 0020-1669  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 124:218506  
 ED Entered STN: 13 Mar 1996  
 AB Zn(II) and Cu(II) complexes of two new potentially pentadentate ligands based on Me 2-aminocyclopent-1-ene-1-dithiocarboxylate with pendant pyrazolyl groups (pzCH<sub>2</sub>)<sub>2</sub>NC<sub>2</sub>H<sub>3</sub>RNHC<sub>5</sub>H<sub>6</sub>CSSCH<sub>3</sub> (R = H, Hmmpcd, and R = CH<sub>3</sub>, Hmmpcd, both having N4S donor atoms set) are reported. The mol. structures of [Zn(hmmpcd)]ClO<sub>4</sub> (1b) and [Cu(hmmpcd)]ClO<sub>4</sub> (2b) show a distorted trigonal bipyramidal geometry for the Zn(II) ion and a square pyramidal geometry for the Cu(II) ion. 1b crystallizes in the triclinic space group P.hivin.1, a 9.900(3), b 15.379(5), c 8.858(2) Å, α 99.93(2), β 93.62(2), γ 100.38(2)°, and Z = 2; while 2b crystallizes in the monoclinic space group P21/n, a 12.859(6), b 12.642(3), c 16.503(2) Å, β 102.67(2)°, and Z = 4. The structures were refined to final R = 0.042 for 1b and 0.049 for 2b. The EPR and electronic spectroscopic studies showed that the Cu(II) species doped into Zn(II) complex adopts the Zn(II) trigonal bipyramidal structure. The cyclic voltammetric measurements indicated 1-electron reversible reduction of the Cu(II) complex occurring at -0.74 V, while irreversible oxidation to Cu(III) takes place at +0.75 V (vs. Ag/AgNO<sub>3</sub>).  
 CC 78-7 (Inorganic Chemicals and Reactions)  
 Section cross-reference(s): 72, 75  
 IT 174280-57-0P 174280-58-1P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and complexation with copper and zinc)  
 IT 174280-57-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and complexation with copper and zinc)  
 RN 174280-57-0 HCAPLUS  
 CN 1-Cyclopentene-1-carbodithioic acid,  
 2-[[2-bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]amino]ethyl]amino]-,  
 methyl ester (CA INDEX NAME)



L47 ANSWER 8 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1996:358780 HCAPLUS Full-text  
 DOCUMENT NUMBER: 125:97176  
 ORIGINAL REFERENCE NO.: 125:18091a,18094a

TITLE: Formation of dinuclear copper(II) complex with  
N,N,N',N'-tetrakis(2-pyridylmethyl)-1,2-ethanediamine  
in aqueous solution

AUTHOR(S): Hirayama, Naoki; Iimuro, Shinji; Kubono, Koji;  
Kokusen, Hisao; Honjo, Takaharu

CORPORATE SOURCE: Dep. Chem., Kanazawa Univ., Kanazawa, 920-11, Japan

SOURCE: Talanta (1996), 43(4), 621-626  
CODEN: TLNTA2; ISSN: 0039-9140

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 20 Jun 1996

AB The polynuclear complexation of divalent 3d transition metal cations with  
N,N,N',N'-tetrakis(2-pyridylmethyl)-1,2-ethanediamine (tpen) in aqueous  
solution was investigated. It was found that copper(II) forms a dinuclear  
complex with tpen in an aqueous solution containing chloride. The composition  
of the complex was determined as Cu<sub>2</sub>Cl<sub>2</sub>(tpen)<sub>2</sub><sup>2+</sup>. Furthermore, the stability  
constant of the complex was determined and its structure was postulated to be  
(μ-Cl)<sub>2</sub>.


CC 68-3 (Phase Equilibria, Chemical Equilibria, and Solutions)

IT 7440-50-8D, Copper, (pyridyl)ethanediamine derivs. complexes  
85264-42-2D, copper complexes 85264-43-3D, copper  
complexes  
RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation,  
nonpreparative)  
(formation of copper(II) complexes with (pyridyl)ethanediamines)

IT 85264-42-2D, copper complexes 85264-43-3D, copper  
complexes  
RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation,  
nonpreparative)  
(formation of copper(II) complexes with (pyridyl)ethanediamines)

RN 85264-42-2 HCAPLUS

CN 1,2-Ethanediamine, N1,N1,N2,N2-tetrakis(1H-pyrazol-1-ylmethyl)- (CA INDEX  
NAME)



RN 85264-43-3 HCAPLUS

CN 1,2-Ethanediamine, N1,N1,N2,N2-tetrakis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]- (CA INDEX NAME)



of 1.97(1) and 1.98(1) Å, while each Zn<sup>2+</sup> ion is further bound to two chloride ions at 2.276(4) and 2.261(4) Å. The Zn<sup>2+</sup> ions are in an almost regular N<sub>2</sub>Cl<sub>2</sub> tetrahedron.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 75

IT 85264-39-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and reactions with zinc chloride and bromide)


IT 85264-39-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and reactions with zinc chloride and bromide)

RN 85264-39-7 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-N1,N2-dimethyl- (CA INDEX NAME)



L47 ANSWER 10 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1994:231747 HCAPLUS Full-text

DOCUMENT NUMBER: 120:231747

ORIGINAL REFERENCE NO.: 120:40821a,40824a

TITLE: Processing of silver halide color photographic material for stable images

INVENTOR(S): Fujita, Yoshihiro; Nakamura, Shigeru

PATENT ASSIGNEE(S): Fuji Photo Film Co Ltd, Japan

SOURCE: Jpn. Kokai Tokyo Koho, 194 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
JP 05034888	A	19930212	JP 1991-214219	19910801 <--
PRIORITY APPLN. INFO.:			JP 1991-214219	19910801 <--


ED Entered STN: 30 Apr 1994

GI For diagram(s), see printed CA Issue.

AB Ag halide color photog. material containing yellow coupler I(R<sub>1</sub> = tertiary alkyl or aryl; R<sub>2</sub> = H, halo, alkoxy, aryloxy, alkyl, dialkylamino; R<sub>3</sub> = substituent; X = heterocyclyl bonded to the coupling active position through aryloxy or N, and releasable on coupling reaction with oxidized aromatic primary amine developer; a = 0-4; R<sub>3</sub> may be same or different when a ≥ 2) is processed by employing a processing solution containing II or III(X = nonmetallic atoms required to form 4-8-membered ring; atoms bonded to N are selected from C, O, S; X<sub>0</sub> = nonmetallic atoms required to form N-containing heterocycle; R<sub>a</sub>, R<sub>b</sub> = alkyl, alkenyl, or bonded to each other to form a 4-8-

membered ring). The processing improved yellow image fastness leading to improved image storage stability.

IC ICM G03C011-00  
ICS G03C007-36  
CC 74-2 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)  
IT 288-13-1, 1H-Pyrazole 288-88-0, 1H-1,2,4-Triazole 76686-85-6  
91272-91-2 91272-92-3 144986-76-5 149310-10-1 150704-12-4  
RL: USES (Uses)  
(yellow image stabilizer, processing solution containing)  
IT 150704-12-4  
RL: USES (Uses)  
(yellow image stabilizer, processing solution containing)  
RN 150704-12-4 HCAPLUS  
CN 1,2-Ethanediamine, N1-ethyl-N2,N2-dimethyl-N1-(1H-pyrazol-1-ylmethyl)-  
(CA INDEX NAME)



L47 ANSWER 11 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1994:123377 HCAPLUS Full-text

DOCUMENT NUMBER: 120:123377

ORIGINAL REFERENCE NO.: 120:21521a,21524a

TITLE: Copper(II) coordination chemistry of potentially octadentate (N8) tetrapyridyl and tetrapyrazolyl-pyridazine ligands. X-ray crystal structures of [Cu<sub>2</sub>(PTAPY)Br<sub>4</sub>].2DMF and [Cu<sub>2</sub>(PTAPY)(NO<sub>3</sub>)<sub>2</sub>(N<sub>3</sub>)(H<sub>2</sub>O)]<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub>.1.2CH<sub>3</sub>OH

AUTHOR(S): Tandon, Santokh S.; Chen, Liqin; Thompson, Laurence K.; Connors, Sean P.; Bridson, John N.

CORPORATE SOURCE: Department of Chemistry, Memorial University of Newfoundland, St. John's, Nfld., A1B 3X7, Can.

SOURCE: Inorganica Chimica Acta (1993), 213(1-2), 289-300

CODEN: ICHAA3; ISSN: 0020-1693

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 05 Mar 1994

AB The structural, electrochem., ESR and magnetic properties of dinuclear Cu(II) complexes of 2 new polyfunctional pyridazine ligands, 3,6-bis(N,N,N',N'-tetrakis(pyridin-2-ylmethyl)aminoethanethiolato)pyridazine (PTAPY), and 3,6-bis(N,N,N',N'-tetrakis(pyrazol-1-ylmethyl)aminoethanethiolato)pyridazine (PTAPZ) are discussed. PTAPY and PTAPZ are potentially octadentate (N8) ligands and on reaction with Cu(II)salts form dinuclear complexes, [Cu<sub>2</sub>(PTAPY)X<sub>4</sub>].yH<sub>2</sub>O (X = NO<sub>3</sub>, y = 1; X = Br, y = 2), [Cu<sub>2</sub>(PTAPY)(MeCN)<sub>2</sub>](ClO<sub>4</sub>)<sub>4</sub>.0.5EtOH, [Cu<sub>2</sub>(PTAPZ)Cl<sub>4</sub>].5H<sub>2</sub>O, and tetranuclear [Cu<sub>2</sub>(PTAPY)(NO<sub>3</sub>)<sub>2</sub>(N<sub>3</sub>)(H<sub>2</sub>O)]<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub>.1.2CH<sub>3</sub>OH (I). The single crystal x-ray structures of [Cu<sub>2</sub>(PTAPY)Br<sub>4</sub>].2DMF (II) and I were determined, and in each case the ligand is hexadentate. Two different 5-coordinate (CuN<sub>3</sub>Br<sub>2</sub>) geometries exist within II. In I, the 2 Cu atoms are quite different with 6- (axially elongated, distorted tetragonal) and 5- (distorted square-pyramidal) coordinate arrangements, and dimerization through azido N atoms gave a

tetranuclear N3- bridged mol. The pyridazine N atoms remain uncoordinated in all complexes and there is no magnetic interaction between the distant Cu centers. Cyclic voltammograms are characterized by the presence of either 1 2-electron or 2 1-electron (overlapping) reversible or quasi-reversible redox processes, associated with the reduction of the dinuclear Cu(II) species to dinuclear Cu(I) species.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 28, 75

IT 152305-82-3P ~~152305-83-4P~~

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and reaction of, with copper salts)


IT ~~152305-83-4P~~

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and reaction of, with copper salts)

RN 152305-83-4 HCAPLUS

CN 1H-Pyrazole-1-methanamine, N,N'-[3,6-pyridazinediylbis(thio-2,1-ethanediyl)]bis[N-(1H-pyrazol-1-ylmethyl)- (9CI) (CA INDEX NAME)



L47 ANSWER 12 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1993:613909 HCAPLUS Full-text

DOCUMENT NUMBER: 119:213909

ORIGINAL REFERENCE NO.: 119:37907a,37910a

TITLE: Method for processing color photographic material

INVENTOR(S): Fujimoto, Hiroshi; Fujita, Yoshihiro; Nakamura, Shigeru

PATENT ASSIGNEE(S): Fuji Photo Film Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 89 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
JP 04362945	A	19921215	JP 1991-193593	19910709 <--
PRIORITY APPLN. INFO.:			JP 1991-99721	A1 19910405 <--
ED Entered STN:		13 Nov 1993		
GI For diagram(s), see printed CA Issue.				

AB In the title processing method, the color photog. material contains  $\geq 1$  high b.p. organic solvent described by O:P(OR1)(OR2)(OR3) and I [R1-5 = alkyl, cycloalkyl, aryl; R6 = halo, alkyl, alkoxy, aryloxy, alkoxycarbonyl; a = 0-3] in its red- and/or green-sensitive photog. emulsion layers, and is processed with a solution containing II and/or III [X = non-metallic atoms required to complete a 4- to 8-membered ring; only C, O, and S can bond to N; X0 = non-metallic atoms required to complete a N-containing hetero-aromatic ring; R7,8 = alkyl, alkenyl; R7 and R8 may form a 4- to 8-membered ring] after being color developed. This method improves stability of magenta dye images.

IC ICM G03C011-00  
ICS G03C001-38; G03C007-388; G03C007-42


CC 74-2 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)  
Section cross-reference(s): 41

IT 951-49-5 78758-54-0 91272-92-3 98816-32-1 144986-84-5  
144986-86-7 146456-97-5 146475-24-3 149310-08-7 150704-05-5  
150704-06-6 150704-07-7, 1H-Pyrazole-1,4-dimethanol 150704-08-8  
150704-09-9 150704-10-2 150704-11-3 150704-12-4  
150704-13-5 150704-14-6 150704-15-7 150704-16-8  
RL: USES (Uses)  
(photog. stabilizer containing)

IT 150704-12-4  
RL: USES (Uses)  
(photog. stabilizer containing)

RN 150704-12-4 HCAPLUS

CN 1,2-Ethanediamine, N1-ethyl-N2,N2-dimethyl-N1-(1H-pyrazol-1-ylmethyl)-  
(CA INDEX NAME)



L47 ANSWER 13 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1993:29868 HCAPLUS Full-text

DOCUMENT NUMBER: 118:29868

ORIGINAL REFERENCE NO.: 118:5361a,5364a

TITLE: Processing solution for silver halide color photographic material

INVENTOR(S): Morigaki, Masakazu; Kawamoto, Hiroyuki; Nakamura, Shigeru

PATENT ASSIGNEE(S): Fuji Photo Film Co., Ltd., Japan

SOURCE: Eur. Pat. Appl., 113 pp.  
CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
EP 504609	A2	19920923	EP 1992-102855	19920220 <--
EP 504609	A3	19930303		
EP 504609	B1	19950719		

R: BE, DE, GB

JP 04313753	A	19921105	JP 1991-142708	19910520 <--
JP 2729542	B2	19980318		
US 5449593	A	19950912	US 1992-838963	19920221 <--
US 5576151	A	19961119	US 1995-426671	19950421 <--
PRIORITY APPLN. INFO.:			JP 1991-48679	A 19910222 <--
			JP 1991-142708	A 19910520 <--
			US 1992-838963	A3 19920221 <--

OTHER SOURCE(S): MARPAT 118:29868

ED Entered STN: 24 Jan 1993

GI For diagram(s), see printed CA Issue.

AB A solution for processing a Ag halide color photog. material for producing color images having excellent storage stability contains  $\geq 1$  compound selected from compds. represented by formulas I, II, and III (Z1 = a nonmetallic atomic group bonding to each N atom with a C, O, or S atom to form a 4- to 8-membered ring; R1, R2 = H, alkyl, alkenyl, aryl, acyl, sulfonyl, sulfinyl, hydroxy, acyloxy, acylamino, sulfonamido, ureido, sulfamoylamino, alkoxycarbonylamino, carbamoyl, sulfamoyl, or heterocyclyl; X1-4 = NR4, N, O, S, CR5R6, CR5, CO, or CNR7 where R4-7 or a substituent group; Z2-4 = a nonmetallic atomic group necessary for forming a 4- to 8-membered ring; Y = O or S; R3 = alkyl, alkenyl, aryl, acyl, sulfonyl, sulfinyl, alkoxycarbonyl, carbamoyl, sulfamoyl, oxalyl, or heterocyclyl).

IC ICM G03C007-30

ICS G03C007-407; G03C007-42

CC 74-2 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)

IT 23230-43-5 67685-12-5 75663-96-6 ~~85264-38-6~~ 91272-74-1  
 91272-91-2 91272-92-3 93597-03-6 101213-03-0 144986-76-5  
 144986-77-6 144986-78-7 144986-79-8 144986-80-1 144986-81-2  
 144986-82-3 144986-83-4 144986-84-5 144986-85-6 144986-86-7  
 144986-87-8 144986-88-9 144986-89-0 144986-90-3 144986-91-4

RL: USES (Uses)

(color photog. stabilizing solns. containing)


IT ~~85264-38-6~~

RL: USES (Uses)

(color photog. stabilizing solns. containing)

RN 85264-38-6 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-dimethyl-N1,N2-bis(1H-pyrazol-1-ylmethyl)- (CA INDEX NAME)



L47 ANSWER 14 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1992:206551 HCAPLUS Full-text

DOCUMENT NUMBER: 116:206551

ORIGINAL REFERENCE NO.: 116:34763a,34766a

TITLE: Transition metal complexes of two related pyrazole containing ligands:  
 3,6-dimethyl-1,8-bis(3,5-dimethyl-1-pyrazolyl)-3,6-diazaoctane (ddad) and  
 1,4-bis(2-(3,5-dimethyl-1-pyrazolyl)ethyl)piperazine (bedp). Synthesis, spectroscopy and x-ray structures



AUTHOR(S): Haanstra, W. G.; Driessen, W. L.; De Graaff, R. A. G.; Sebrechts, G. C.; Suriano, J.; Reedijk, J.; Turpeinen, U.; Hamalainen, R.; Wood, J. S.

CORPORATE SOURCE: Dep. Chem., Leiden Univ., Leiden, 2300 RA, Neth.

SOURCE: Inorganica Chimica Acta (1991), 189(2), 243-51  
CODEN: ICHAA3; ISSN: 0020-1693

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 16 May 1992

AB Several coordination compds. with 3,6-dimethyl-1,8-bis(3,5-dimethyl-1-pyrazolyl)-3,6-diazaoctane (ddad) were obtained: [M(ddad)](BF<sub>4</sub>)<sub>2</sub> (I; M = Cu, Ni), Co(ddad)(H<sub>2</sub>O)(BF<sub>4</sub>)<sub>2</sub>, M<sub>2</sub>(ddad)Cl<sub>4</sub> (II; M = Co, Ni, Cu, Zn), Co<sub>3</sub>(ddad)<sub>2</sub>(NCS)<sub>6</sub> (III) and Cu<sub>2</sub>(ddad)(NCS)<sub>3</sub> (IV). Five x-ray structures were obtained, viz. of I, Ni(ddad)(NCS)<sub>2</sub> (V), [Ni(bedp)](H<sub>2</sub>O)(BF<sub>4</sub>)<sub>2</sub> (bedp = 3,6-dimethyl-1,8-bis(3,5-dimethyl-1-pyrazolyl)-3,6-diazaoctane) and [Cu(bedp)](H<sub>2</sub>O)(BF<sub>4</sub>)<sub>2</sub>, but the data did not allow the calcn. of very accurate bond lengths. However, the basic coordination geometries were obtained in all cases. The coordination by the ligands is square planar, while V contains addnl. trans coordinating thiocyanate anions. The asym. unit of I (M = Cu) contains two almost identical [Cu(ddad)]<sup>2+</sup> species. The coordination of the copper atoms in both mols. is intermediate between tetrahedral and square planar. [Cu(ddad)]<sup>2+</sup> exists as the (R,R) and (S,S) diastereoisomers of the ligand. II are, except for M = Ni, all dinuclear with MN<sub>2</sub>Cl<sub>2</sub> chromophores. II (M = Ni) contains square planar Ni(ddad)<sub>2</sub><sup>+</sup> cations and tetrahedral NiCl<sub>4</sub><sup>2-</sup> anions. V, which is isomorphous with the corresponding Zn(II) compds., has octahedral MN<sub>2</sub>N<sub>2</sub>'N<sub>2</sub>' chromophores. III crystallizes as [Co(ddad)(NCS)]<sub>2</sub>Co(NCS)<sub>4</sub> with five coordinate cobalt in the cation. IV is formulated as [Cu(ddad)]<sup>2+</sup> and [Cu(NCS)<sub>3</sub>]<sup>2-</sup>. [M(bedp)](H<sub>2</sub>O)(BF<sub>4</sub>)<sub>2</sub> (M = Ni, Cu), M(bedp)(NCS)<sub>2</sub> (M = Ni, Co), Zn<sub>2</sub>(bedp)(NCS)<sub>4</sub>, M<sub>2</sub>(bedp)Cl<sub>4</sub> (VI; M = Ni, Co, Cu, Zn) and Cu<sub>2</sub>(bedp)(NCS)<sub>3</sub> (VII) were obtained. VI form dinuclear compds. (similar to ddad) with MN<sub>2</sub>Cl<sub>2</sub> chromophores. However, in addition to the green form of [Ni(bedp)][NiCl<sub>4</sub>] a purple isomer Ni<sub>2</sub>(bedp)Cl<sub>4</sub>, with a tetrahedral NiN<sub>2</sub>Cl<sub>2</sub> chromophore, was obtained. The ligand field spectra of V and the isomorphous cobalt compound show typical octahedral chromophores. VII has a structure likely to be similar to the corresponding ddad compound, viz. [Cu(II)(bedp)][Cu(I)(SCN)<sub>3</sub>]. The preps. of the ligands are described.


CC 78-7 (Inorganic Chemicals and Reactions)  
Section cross-reference(s): 28, 75

IT 139775-87-4P 139775-88-5P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and complexation of)

IT 139775-87-4P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and complexation of)

RN 139775-87-4 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-N1,N2-dimethyl- (CA INDEX NAME)



L47 ANSWER 15 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1989:184768 HCAPLUS Full-text

DOCUMENT NUMBER: 110:184768

ORIGINAL REFERENCE NO.: 110:30457a,30460a

TITLE: Dinuclear transition metal compounds of a decadentate pyrazole-containing chelating ligand. X-ray crystal structure of  $\text{Co}_2(\text{tthd})(\text{ClO}_4)_4(\text{H}_2\text{O})_2(\text{MeOH})1.75$ 

AUTHOR(S): Paap, F.; Driessen, W. L.; De Graaff, R. A. G.; Reedijk, J.

CORPORATE SOURCE: Dep. Chem., Leiden Univ., Leiden, 2300 RA, Neth.

SOURCE: Polyhedron (1988), 7(24), 2575-81

CODEN: PLYHDE; ISSN: 0277-5387

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 12 May 1989

AB A potentially decadentate ligand, 1,1,4,7,10,10-hexakis(3,5-dimethyl-1-pyrazolylmethyl)-1,4,7,10-tetraazadecane ( $\text{H}_2\text{tthd}$ ), was prepared from the reaction of triethylenetetramine with 6 equiv of N-hydroxymethyl-3,5-dimethylpyrazole.  $\text{M}_2(\text{tthd})(\text{ClO}_4)_4(\text{H}_2\text{O})_x$  ( $\text{M} = \text{Co}, \text{Ni}, \text{Cu}, \text{Zn}, \text{Cd}; x = 4-8$ ) and  $\text{M}_2(\text{tthd})\text{X}_2(\text{ClO}_4)_2(\text{H}_2\text{O})_x$  ( $\text{M} = \text{Co}, \text{Ni}; \text{X} = \text{NCS}, \text{Cl}; x = 4-8$ ) were prepared.  $\text{Co}_2(\text{tthd})(\text{ClO}_4)_4(\text{H}_2\text{O})_2(\text{MeOH})1.75$  crystallizes in the triclinic space group  $\text{P}1$ ,  $a$  1.959(2),  $b$  1.5657(3),  $c$  2.1244(3) nm,  $\alpha$  105.5(1),  $\beta$  96.9(1),  $\gamma$  112.1(1)°. Due to severe disorder of the anions the structure could only be refined to  $R_w = 0.099$ . The ligand acts as a decadentate, dinucleating ligand. The Co ions are distorted octahedrally, surrounded by 5 N-atoms of the tthd ligand and an O atom of water occupying the 6th coordination place. The other perchlorate compds. have very similar structures, as can be concluded from spectroscopic data. In the thiocyanate and chloride compds. the anions replace coordinated  $\text{H}_2\text{O}$  mols., resulting in octahedral Ni compds. With Co thiocyanate, however, tthd acts as an octadentate ligand, resulting only in 5-coordinated compds.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 28, 75

IT 85264-45-5P 119781-62-3P 120170-22-1P 120170-25-4P  
120170-28-7P 120183-41-7P 120183-44-0P 120293-21-2P


RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

IT 85264-45-5P


RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 85264-45-5 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis[2-[bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]amino]ethyl]-N1,N2-bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-  
(CA INDEX NAME)



L47 ANSWER 16 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1989:185371 HCAPLUS Full-text  
 DOCUMENT NUMBER: 110:185371  
 ORIGINAL REFERENCE NO.: 110:30551a,30554a  
 TITLE: 3,5-Diphenyl-1H-pyrazole derivatives. II.  
 N-Substituted 1-(2-aminoethyl)-3,5-diphenyl-1H-  
 pyrazoles and their 4-bromo derivatives with analgesic  
 and other activities  
 AUTHOR(S): Bondavalli, F.; Bruno, O.; Ranise, A.; Schenone, P.;  
 Russo, S.; Loffreda, A.; De Novellis, V.; Lo Sasso,  
 C.; Marmo, E.  
 CORPORATE SOURCE: Ist. Sci. Farm., Univ. Genova, Genoa, Italy  
 SOURCE: Farmaco, Edizione Scientifica (1988),  
 43(12), 1019-34  
 CODEN: FRPSAX; ISSN: 0430-0920  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 110:185371  
 ED Entered STN: 26 May 1989  
 GI



AB Eighteen title compds. (I, R = H, alkyl; R<sub>1</sub> = alkyl; R<sub>1</sub> = alkylene, alkyleneoxyalkylene, alkyleneiminoalkylene, and etc.; R<sub>2</sub> = H, Br) were prepared by reaction of the corresponding 1-(2-hydroxyethyl) compds. with tosyl chloride, followed by refluxing the resultant tosylates with an excess of primary or secondary amine. I had marked analgesic activity. Moreover, various I had moderate hypotensive, bradycardic, anti-inflammatory, and local anesthetic activity in vivo, as well as weak platelet-antiaggregating potency in vitro.

CC 1-4 (Pharmacology)

Section cross-reference(s): 28

IT 4162-98-5P 120217-55-2P 120217-56-3P 120217-57-4P  
 120217-58-5P 120217-59-6P 120217-60-9P 120217-61-0P 120217-62-1P  
 120217-63-2P 120217-64-3P 120217-65-4P 120217-66-5P 120217-67-6P  
 120217-68-7P 120217-69-8DP, derivs. 120217-72-3P 120217-73-4P  
 120217-74-5P 120217-75-6P 120217-76-7P 120217-77-8P 120217-78-9P  
 120217-79-0P 120217-80-3P 120217-81-4P 120217-82-5P 120217-83-6P  
 120253-58-9P 120253-59-0P 120253-60-3P

RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)  
 (preparation and pharmacol. of)

IT 120217-57-4P 120217-73-4P


RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic

10/551,292

preparation); THU (Therapeutic use); BIOL (Biological study); PREP  
(Preparation); PROC (Process); USES (Uses)  
(preparation and pharmacol. of)


RN 120217-57-4 HCAPLUS

CN 1,2-Ethanediamine, N'-[2-(3,5-diphenyl-1H-pyrazol-1-yl)ethyl]-N,N-dimethyl-  
(9CI) (CA INDEX NAME)



RN 120217-73-4 HCAPLUS

CN 1,2-Ethanediamine, N'-[2-(3,5-diphenyl-1H-pyrazol-1-yl)ethyl]-N,N-dimethyl-  
, dihydrochloride (9CI) (CA INDEX NAME)



●2 HCl

L47 ANSWER 17 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1988:197156 HCAPLUS Full-text

DOCUMENT NUMBER: 108:197156

ORIGINAL REFERENCE NO.: 108:32217a,32220a

TITLE: Copper complexes of some tetradentate pyrazolyl amines

AUTHOR(S): Addison, A. W.; Palaniandavar, M.; Driessen, W. L.;  
Paap, F.; Reedijk, J.

CORPORATE SOURCE: Chem. Dep., Drexel Univ., Philadelphia, PA, 19104, USA

SOURCE: Inorganica Chimica Acta (1988), 142(1),  
95-100


CODEN: ICHAA3; ISSN: 0020-1693

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 28 May 1988

GI



I

AB I [R = R1 = H (edbp); R = Me, R1 = H (edbd); R = H, R1 = Me (debp); R = R1 = Me (debd)] were prepared by condensation of N-hydroxymethylpyrazoles with ethylenediamines. CuL(ClO4)2 (L = debp, edbp, edbd) and Cu(debd)(dMp)X2 (X = ClO4, BF4; dMp = 3,5-dimethylpyrazole) were prepared. The compds. were characterized by their absorption (d-d) and EPR spectra. All 4 undergo quasi-reversible electrochem. reduction in MeOH, the redox potentials being correlated with the degree of ligand methylation. The Cu(I) complexes are relatively unstable and bind CO with different affinities.


CC 78-7 (Inorganic Chemicals and Reactions)

IT 114287-01-3P 114315-22-9P 114315-23-0P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)

IT 114315-22-9P 114315-23-0P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)


RN 114315-22-9 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis(1H-pyrazol-1-ylmethyl)- (CA INDEX NAME)



RN 114315-23-0 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]- (CA INDEX NAME)



L47 ANSWER 18 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1989:106933 HCAPLUS Full-text

DOCUMENT NUMBER: 110:106933

ORIGINAL REFERENCE NO.: 110:17483a,17486a

TITLE: Transition metal complexes of ligands containing 3,5-dimethylpyrazolyl groups and thioether functions.

The crystal and molecular structure of (1,12-bis(3,5-dimethylpyrazol-1-yl)-2,11-diaza-5,8-

dithiadodecane)nickel(II) bis(tetrafluoroborate)

AUTHOR(S): Paap, F.; Driessen, W. L.; Reedijk, J.; Spek, A. L.

CORPORATE SOURCE: Dep. Chem., Leiden Univ., Leiden, 2300 RA, Neth.

SOURCE: Inorganica Chimica Acta (1988), 150(1), 57-64

CODEN: ICHAA3; ISSN: 0020-1693

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 17 Mar 1989

AB The hexadentate (N4S2) ligand 1,12-bis(3,5-dimethylpyrazol-1-yl)-2,11-diaza-5,8-dithiadodecane (dsbd) forms M(dsbd)A2 (M = Co, Ni or Cd; A = ClO4 or BF4).

Ni(dsbd)(BF<sub>4</sub>)<sub>2</sub> crystallizes in the monoclinic space group P2<sub>1</sub>/n, Z = 4, a 1.2817(4), b 1.5512(3), c 1.33666(4) nm, β 94.11(1)°, R<sub>w</sub> = 0.032 for 2181 observed reflections. The Ni ion is octahedrally chelated by 2 S atoms, 2 amine nitrogens and 2 pyrazole nitrogens. Ni-N amts. to 210-212 pm, Ni-S distances are 243.5 pm. The other dsbd compds. have very similar structures as is concluded from their spectroscopic properties. The octadentate (N<sub>6</sub>S<sub>2</sub>) ligand 1,1,10,10-tetrakis(3,5-dimethylpyrazol-1-ylmethyl)-1,10-diaza-4,7-dithiadecane (dstd) forms M<sub>2</sub>(dstd)A<sub>4</sub>(H<sub>2</sub>O)<sub>n</sub>, (M = Mn, Fe, Co, Ni, Cu, Zn or Cd; A = ClO<sub>4</sub>, n = 4; M = Co, Ni or Zn, A = BF<sub>4</sub>, n = 4; M = Co or Ni, A = NCS, n = 0). In all compds. the metal ions are 6-coordinated, as deduced from their spectroscopic properties.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 75

IT 118202-79-2P, 1,1,10,10-Tetrakis(3,5-dimethylpyrazol-1-ylmethyl)-1,10-diaza-4,7-dithiadecane 118202-80-5P,

1,12-Bis(3,5-dimethylpyrazol-1-yl)-2,11-diaza-5,8-dithiadodecane

118473-35-1P 118473-36-2P 118473-38-4P 118473-39-5P 118473-42-0P

118473-43-1P 118473-45-3P 118504-38-4P 118504-40-8P 118504-41-9P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)

IT 118202-79-2P, 1,1,10,10-Tetrakis(3,5-dimethylpyrazol-1-ylmethyl)-1,10-diaza-4,7-dithiadecane 118202-80-5P,

1,12-Bis(3,5-dimethylpyrazol-1-yl)-2,11-diaza-5,8-dithiadodecane


RL: SPN (Synthetic preparation); PREP (Preparation)


(preparation of)

RN 118202-79-2 HCAPLUS

CN 1H-Pyrazole-1-methanamine, N,N'-[1,2-ethanediylbis(thio-2,1-ethanediyl)]bis[N-[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-3,5-dimethyl-  
(9CI) (CA INDEX NAME)



PAGE 1-A





RN 118202-80-5 HCAPLUS

CN 1H-Pyrazole-1-methanamine, N,N'-[1,2-ethanediylbis(thio-2,1-ethanediyl)]bis[3,5-dimethyl- (9CI) (CA INDEX NAME)



L47 ANSWER 19 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1987:94880 HCAPLUS Full-text

DOCUMENT NUMBER: 106:94880

ORIGINAL REFERENCE NO.: 106:15371a,15374a

TITLE: Five- and six-coordinated transition-metal mixed-anion compounds containing the sterically constrained pyrazole-containing ligand 1,6-bis(3',5'-dimethylpyrazol-1'-yl)-2,5-dimethyl-2,5-diazahexane

AUTHOR(S): Paap, Frans; Driessen, Willem L.; Reedijk, Jan

CORPORATE SOURCE: Dep. Chem., State Univ. Leiden, Leiden, 2300 RA, Neth.

SOURCE: Polyhedron (~~1986~~), 5(11), 1815-19

CODEN: PLYHDE; ISSN: 0277-5387

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 21 Mar 1987

AB The dimethylpyrazole derivative of N,N-dimethylethylenediamine, viz. 1,6-bis(3',5'-dimethylpyrazol-1'-yl)-2,5-diazahexane (debd), forms coordination compds. with the transition-metal ions (M = Co(II), Ni(II), Cu(II) and Zn(II)) in the presence of a halide and a ClO4--anion of stoichiometry [M(debd)X(H2O)n]ClO4 (n = 0 or 1). The ligand is always tetradentate. With Co and Ni 6-coordinate compds. are formed. With Cu and Zn 5-coordinate

compds. are formed. The compds. were characterized by spectroscopic and magnetic methods (IR, ligand field, ESR and NMR).

CC 78-7 (Inorganic Chemicals and Reactions)

IT 85264-39-7, 1,6-Bis(3,5-dimethylpyrazol-1-yl)-2,5-dimethyl-2,5-diazahehexane


RL: PRP (Properties)  
(NMR of)

IT 85264-39-7, 1,6-Bis(3,5-dimethylpyrazol-1-yl)-2,5-dimethyl-2,5-diazahehexane

RL: PRP (Properties)  
(NMR of)

RN 85264-39-7 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-N1,N2-dimethyl- (CA INDEX NAME)



L47 ANSWER 20 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1986:140923 HCAPLUS Full-text

DOCUMENT NUMBER: 104:140923

ORIGINAL REFERENCE NO.: 104:22097a,22100a

TITLE: Copper complexes with quadridentate bis(pyrazolyl)thioether amine and tris(pyrazolyl)amine ligands. Structural characterization of the complexes isothiocyanato[tris(2-pyrazolylethyl)amine-NN2N2'N2'']copper(II) diisothiocyanatocuprate(I) and {bis[2-(3',5'-dimethylpyrazolyl)ethyl](2-methylthioethyl)amine-NN2N2'S}chlorocopper(II) chloride dihydrate

AUTHOR(S): Di Vaira, Massimo; Mani, Fabrizio

CORPORATE SOURCE: Dep. Chem., Univ. Florence, Florence, Italy

SOURCE: Journal of the Chemical Society, Dalton Transactions: Inorganic Chemistry (1972-1999) (1985), (11), 2327-32

CODEN: JCDBTBI; ISSN: 0300-9246

DOCUMENT TYPE: Journal

LANGUAGE: English


ED Entered STN: 19 Apr 1986

AB Reactions of simple Cu salts with the tripod quadridentate ligands L [L = tris(2-pyrazolylethyl)amine (L1), bis[2-(3',5'-dimethylpyrazolyl)ethyl][2-(methylthio)ethyl]amine (L2)] gave the complexes [CuXL1][BPh4] (X = Cl, Br, NCS), [CuClL1]2[CuCl4], [Cu(NCS)L][Cu(NCS)2] (I), [Cu(NCS)L2][BPh4].Me2CO, [CuClL2]Cl.2H2O (II), and [CuBrL2]3[CuBr3]Br.H2O. The crystal and mol. structures of I (L = L1) and II were determined by x-ray diffraction using the heavy-atom method and refined by full-matrix least squares to R 0.040 and 0.044 for 2518 and 1559 observed reflections, resp. I (L = L1) consists of isolated cations with the Cu in an approx. trigonal-bipyramidal environment of 5 N atoms and of layers formed by the [Cu(NCS)2]- anion, with pseudotetrahedral Cu(I) atoms. The Cu(II) atom in the cation of II is in a square-pyramidal environment of 1 S, 1 Cl, and 3 N atoms. L2 was prepared by the reaction of bis(2-chloroethyl)amine (which causes blistering) with K 3,5-




dimethylpyrazolate in THF, followed by reaction of the intermediate amine with Cl(CH<sub>2</sub>)<sub>2</sub>SMe in EtOH.

CC 78-7 (Inorganic Chemicals and Reactions)  
 Section cross-reference(s): 28, 75  
 IT 101049-76-7P 101049-79-0P 101049-81-4P 101065-14-9P 101065-15-0P  
 101065-18-3P 101125-00-2P 101131-44-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)  
 IT 101131-44-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)  
 RN 101131-44-6 HCAPLUS  
 CN 1H-Pyrazole-1-ethanamine, N-[2-(methylthio)ethyl]-N-[2-(1H-pyrazol-1-yl)ethyl]- (CA INDEX NAME)




L47 ANSWER 21 OF 23 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1984:530693 HCAPLUS Full-text  
 DOCUMENT NUMBER: 101:130693  
 ORIGINAL REFERENCE NO.: 101:19885a,19888a  
 TITLE: (Azolylmethyl)amines and their use in microbicidal agents  
 INVENTOR(S): Oeckl, Siegfried; Schmitt, Hans Georg; Paulus, Wilfried; Genth, Hermann  
 PATENT ASSIGNEE(S): Bayer A.-G. , Fed. Rep. Ger.  
 SOURCE: Ger. Offen., 38 pp.  
 CODEN: GWXXBX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3238006	A1	19840419	DE 1982-3238006	19821013 <--
US 4599427	A	19860708	US 1983-536929	19830928 <--
EP 106243	A1	19840425	EP 1983-109794	19830930 <--
EP 106243	B1	19860528		
R: BE, DE, FR, GB, IT, SE				
JP 59093058	A	19840529	JP 1983-188558	19831011 <--
JP 06029268	B	19940420		
CA 1243315	A1	19881018	CA 1983-438858	19831011 <--
PRIORITY APPLN. INFO.:			DE 1982-3238006	A 19821013 <--
OTHER SOURCE(S): CASREACT 101:130693				
ED Entered STN: 13 Oct 1984				
GI				



- AB R1R2NCH2R (R = Q, Q1; X = N, CR7; R1 = C1-24 aliphatic group, C3-12 cycloalkyl, C7-24 aralkyl, C1-24 aminoalkyl, C2-24 alkylenaminoalkyl, C6-18 aryl, C7-24 alkylaryl, C6-18 haloaryl, C1-24 alkoxy, C7-24 arylalkoxy, CH2R; R2 = H, R1; NR1R2 = 5- or 6-membered heterocyclcyl; R3-R7 = H, halo, NO2, C1-24 alkyl, C3-12 cycloalkyl, C1-24 alkoxy, cyano, C7-24 aralkyl), useful as bactericides, fungicides, algicides, and slimicides, were prepared Thus, treating [Me(CH2)7]2NH and 1,2,4-triazole in CH2Cl2 or (ClCH2)2 with 30% HCHO at 30-35° gave 98% (triazolylmethyl)amine I. The min. inhibitory concentration of I for Alternaria tenuis was 50 µg/L, for Escherichia coli was 95 µg/L, for slime organisms 75 µg/L, and for a mixed culture of green, blue, and brown algae and diatoms the lethal concentration was 20 µg/L.
- IC C07D249-08; C07D275-04; C07D401-06; C07D403-06; C07D413-06; A01N043-50; A01N043-64; A01N043-80; A01N043-00; C02F001-50; C09D005-14; C09J003-00
- CC 28-10 (Heterocyclic Compounds (More Than One Hetero Atom))  
Section cross-reference(s): 5, 10
- IT 23230-39-9P 23230-41-3P 23230-42-4P 23230-43-5P 23230-44-6P  
76686-85-6P 76686-86-7P 91272-74-1P 91272-75-2P 91272-76-3P  
91272-77-4P 91272-78-5P 91272-79-6P 91272-80-9P 91272-81-0P  
91272-82-1P 91272-83-2P 91272-84-3P 91272-85-4P 91272-86-5P  
91272-87-6P 91272-88-7P 91272-89-8P 91272-90-1P 91272-91-2P  
91272-92-3P 91272-93-4P 91272-94-5P 91272-95-6P 91272-96-7P  
91272-97-8P 91272-98-9P 91272-99-0P 91273-00-6P 91273-01-7P  
91273-08-4P 91273-09-5P 91273-10-8P 91273-11-9P  
91273-12-0P 91273-13-1P 91273-14-2P 91273-15-3P 91273-16-4P  
91273-17-5P 91273-18-6P 91273-19-7P 91273-20-0P 91273-22-2P  
91273-23-3P 91273-24-4P 91273-26-6P 91273-30-2P 91273-31-3P  
91273-32-4P 91273-33-5P 91273-34-6P 91273-35-7P 91273-36-8P  
91273-37-9P 91273-39-1P 91274-94-1P 93597-03-6P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of, as biocide)
- IT 91273-11-9P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of, as biocide)
- RN 91273-11-9 HCAPLUS
- CN 1,2-Ethanediamine, N1,N2-dimethyl-N1-[2-[methyl(1H-pyrazol-1-ylmethyl)amino]ethyl]-N2-(1H-pyrazol-1-ylmethyl)- (CA INDEX NAME)



ORIGINAL REFERENCE NO.: 98:24379a,24382a  
TITLE: Synthesis of some new pyrazole-containing  
chelating agents  
AUTHOR(S): Driessen, Willem L.  
CORPORATE SOURCE: Dep. Chem., State Univ. Leiden, Leiden, 2300 RA, Neth.  
SOURCE: Recueil: Journal of the Royal Netherlands Chemical  
Society (1982), 101(12), 441-3  
CODEN: RJRSDK; ISSN: 0165-0513  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 98:160629  
ED Entered STN: 12 May 1984  
GI



AB The (hydroxymethyl)pyrrazoles I (R = H, Me) underwent condensation with EtNH<sub>2</sub>, PhNH<sub>2</sub>, MeNHCH<sub>2</sub>CH<sub>2</sub>NHMe, H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, (H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>NHCH<sub>2</sub>)<sub>2</sub>, and NH<sub>3</sub> to give products in which all amine hydrogens were substituted by 1-pyrrazolylmethyl groups, e.g. II.

CC 28-8 (Heterocyclic Compounds (More Than One Hetero Atom))

ST chelating agent pyrazolylmethylanine deriv; pyrazolylmethylanine  
deriv

IT	85264-34-2P	85264-35-3P	85264-36-4P	85264-37-5P	<del>85264-38-6P</del>
	<del>85264-39-7P</del>	85264-40-0P	85264-41-1P	<del>85264-42-2P</del>	
	<del>85264-43-3P</del>	<del>85264-44-4P</del>	<del>85264-45-5P</del>		
	<del>85264-46-6P</del>	<del>85264-47-7P</del>	85264-48-8P	85264-49-9P	


RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

IT	<u>85264-38-6P</u>	<u>85264-39-7P</u>	<u>85264-42-2P</u>
	<u>85264-43-3P</u>	<u>85264-44-4P</u>	<u>85264-45-5P</u>
	<u>85264-46-6P</u>	<u>85264-47-7P</u>	

RL: SPN (Synthetic preparation); PREP (Preparation  
(preparation of))


RN 85264-38-6 HCAPLUS

CN	1,2-Ethanediamine, N1,N2-dimethyl-N1,N2-bis(1H-pyrazol-1-ylmethyl)-	(CA
	INDEX NAME)	




RN 85264-39-7 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-N1,N2-dimethyl- (CA INDEX NAME)




RN 85264-42-2 HCAPLUS

CN 1,2-Ethanediamine, N1,N1,N2,N2-tetrakis(1H-pyrazol-1-ylmethyl)- (CA INDEX NAME)



RN 85264-43-3 HCAPLUS


CN 1,2-Ethanediamine, N1,N1,N2,N2-tetrakis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]- (CA INDEX NAME)



RN 85264-44-4 HCAPLUS


CN 1,2-Ethanediamine, N1,N2-bis[2-[bis(1H-pyrazol-1-ylmethyl)amino]ethyl]-N1,N2-bis(1H-pyrazol-1-ylmethyl)- (CA INDEX NAME)

10/551,292




RN 85264-45-5 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis[2-[bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]amino]ethyl]-N1,N2-bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]- (CA INDEX NAME)




RN 85264-46-6 HCAPLUS

CN 1,2-Ethanediamine, N1-[2-[bis(1H-pyrazol-1-ylmethyl)amino]ethyl]-N1,N2,N2-tris(1H-pyrazol-1-ylmethyl)- (CA INDEX NAME)



RN 85264-47-7 HCAPLUS

CN 1,2-Ethanediamine, N1-[2-[bis[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]amino]ethyl]-N1,N2,N2-tris[(3,5-dimethyl-1H-pyrazol-1-yl)methyl]- (CA INDEX NAME)



=> diall abeq tech abex fraghitstr 23

YOU HAVE REQUESTED DATA FROM FILE 'HCAPLUS, WPIX' - CONTINUE? (Y)/N:y

L47 ANSWER 23 OF 23 WPIX COPYRIGHT 2009 THOMSON REUTERS on STN  
 ACCESSION NUMBER: 2004-068858 [07] WPIX  
 CROSS REFERENCE: 1996-221899; 1998-130312; 1998-240752; 1998-240943;  
 1998-240950; 1998-241072; 1998-397917; 1998-583442;  
 1999-080876; 1999-132432; 1999-142839; 1999-142962;  
 1999-302357; 1999-611065; 2000-052183; 2000-105073;  
 2000-136305; 2000-338767; 2000-505663; 2000-638498;  
 2001-272561; 2001-272673; 2001-391546; 2001-656196;  
 2002-061387; 2002-138760; 2002-156532; 2002-314807;  
 2002-463278; 2002-470122; 2002-625845; 2003-038162;  
 2003-329825; 2003-354490; 2003-656115; 2003-669409;  
 2003-742808; 2003-754498; 2003-765734; 2003-777476;  
 2003-811094; 2004-212114; 2004-223799; 2004-224691;  
 2004-236784; 2006-008320; 2008-C33816  
 DOC. NO. CPI: C2004-028387 [07]  
 DOC. NO. NON-CPI: N2004-055373 [07]  
 TITLE: Production of metal-ligand compounds used in olefin  
 oligomerization, by synthesizing first and second metal  
 binding ligands in respective regions on substrate and  
 delivering respective metal ion to each metal binding  
 ligand  
 DERWENT CLASS: A60; B04; E19; J04; S03  
 INVENTOR: BOUSSIE T; GOLDWASSER I; MCFARLAND E; MURPHY V; POWERS T;  
 TURNER H; VAN BEEK J A M; WEINBERG W H  
 PATENT ASSIGNEE: (SYMY-N) SYMYX TECHNOLOGIES INC  
 COUNTRY COUNT: 1

PATENT INFORMATION:

PATENT NO	KIND	DATE	WEEK	LA	PG	MAIN IPC
US 20030100119	A1	20030529	(200407)*	EN	64	[30]

<--

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
-----------	------	-------------	------

US 20030100119 A1 Div Ex	US 1994-327513 19941018
US 20030100119 A1 Provisional	US 1996-16102P 19960723
US 20030100119 A1 Provisional	US 1996-28106P 19961009
US 20030100119 A1 Provisional	US 1996-29255P 19961025
US 20030100119 A1 Provisional	US 1997-35366P 19970110
US 20030100119 A1 Provisional	US 1997-48987P 19970609
US 20030100119 A1 CIP of	US 1998-127660 19980731
US 20030100119 A1 Cont of	US 1999-337047 19990621
US 20030100119 A1	US 2002-269362 20021011

## FILING DETAILS:

PATENT NO	KIND	PATENT NO
US 20030100119 A1	Div ex	US 5985356 A
US 20030100119 A1	CIP of	US 6420179 B

PRIORITY APPLN. INFO: US 2002-269362 20021011

<u>US 1994-327513</u>	<u>19941018</u>
<u>US 1996-16102P</u>	<u>19960723</u>
<u>US 1996-28106P</u>	<u>19961009</u>
<u>US 1996-29255P</u>	<u>19961025</u>
<u>US 1997-35366P</u>	<u>19970110</u>
<u>US 1997-48987P</u>	<u>19970609</u>
<u>US 1998-127660</u>	<u>19980731</u>
<u>US 1999-337047</u>	<u>19990621</u>

## INT. PATENT CLASSIF.:

IPC RECLASSIF.: G01N0001-10 [I,A]; G01N0001-10 [I,C]  
 USCLASS NCLM: 436/037.000  
 NCLS: 436/073.000; 436/180.000

## BASIC ABSTRACT:

US 20030100119 A1 UPAB: 20060203

NOVELTY - Production of an array of metal ligand compounds (A) comprises synthesizing first and second metal binding ligands in respective regions on a substrate and delivering respective metal ions to each metal binding ligand to form first and second metal ligand compounds.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for:

(a) preparing a polymer blend by contacting at least two (A) with a cocatalyst and a monomer, and

(b) polymerizing olefins, diolefins, and acetylenically unsaturated monomers by contacting (A) with a cocatalyst and a monomer.

USE - Used for preparing an array of (A) useful for an organic transformation reaction requiring Lewis acidic sites, e.g. stereo-selective coupling reactions, olefin oligomerization reactions or olefin polymerization reactions. (A) Are used to prepare polymer blend or to polymerize olefins, diolefins or acetylenically unsaturated monomers.

ADVANTAGE - The method accelerates the rate discovering and optimizing catalytic process, and rapidly characterizes each member to identify compounds with specific, desired properties, e.g. polymerization characteristics, mechanical, optical, physical or morphological property, lifetime, stability, selectivity, conversion efficiency, or activity of (A).

MANUAL CODE: CPI: A02-A06D; A04-G01A; B05-A03B; B07-D08; B10-A20;  
 B11-C01A; B11-C01B; E05-L02C; E05-M; E11-K01; E11-K02;  
 J04-E04  
 EPI: S03-E13D

## TECH

INORGANIC CHEMISTRY - Preferred Method: The method also involves activating both metal-ligands with respective activator to form first and second activated (A). The synthesizing step also involves sequentially

synthesizing first and second components of each metal binding ligand. The method also involves screening the array of (A) for useful properties using scanned mass spectrometry, chromatography, ultraviolet imaging, visible imaging, infrared imaging, electromagnetic imaging, ultraviolet spectroscopy, visible spectroscopy, infrared spectroscopy, electromagnetic spectroscopy, or acoustical methods.

The array of (A) is further modified by reaction with an ion-exchange activator to produce an array of ligand-stabilized cationic aluminum reagents.

Preferred Components: The activated (A) are organometallic compounds, homogeneous catalysts or heterogeneous catalysts. The first and second metal-ligand compounds are activated-free catalysts, which are homogeneous or heterogeneous catalysts. The metal-binding ligands are neutral bidentate ligands, mono-anionic bidentate ligands, chelating diamine ligands comprising 1,2-diamine ligands, salen ligands or ancillary ligands. The first and second metal ions are each transition metals comprising palladium, nickel, platinum, iridium, rhodium, chromium, molybdenum, tungsten, or cobalt. The first and second activators include methylaluminoxane (MAO), (Q)+(NCA)- (NCA = non-coordinating anions), (H(OEt<sub>2</sub>)+(BAr<sub>4</sub>)- or (H(OEt<sub>2</sub>))+(B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>)-.

The metal-binding ligands are (2,2) or (2,1) ligands that are respectively contacted with a main group of metal alkyl complex such that they are in the mono- or di-protic form. The ligand-stabilized cationic aluminum reagents can be used as catalyst for the transformation reaction.

The metal-binding ligands have a coordination number (CN) comprising 1-3 or a charge comprising 0, -1 to -4. They have a CN with corresponding charge consisting of CN = 2, charge = -2, -1, -3 or neutral; CN = 1, charge = -1, -2 or -3; or CN = 3, charge = -1 or -3.

When metal-binding ligands are ancillary ligands, each has a charge that is greater than the coordination number, neutral bidentate ligands, each of the transition metal ions is stabilized by a labile neutral Lewis base, chelating diamine ligands each of the transition metal ions is a Group 10 transition metal, or monoanionic bidentate ligands, each of the transition metal ions is stabilized by a labile anionic leaving group ligand. The activators form counter-ions after activation. The metal-binding ligands are directly attached to the substrate or to the first and second synthesis support on the substrate, or attached to the substrate through first and second linker groups. The substrate is a porous or non-porous substrate. The array can be screened simultaneously, serially, or in a spatially selective manner.

Each (A) is synthesized in an area of less than 25 cm<sup>2</sup>, preferably less than 1 mm<sup>2</sup>, especially less than 1 micro-m<sup>2</sup>. At least 10 (especially at least 10 power 6) different (A) are synthesized on the substrate.

ORGANIC CHEMISTRY - Preferred Components: The metal alkyl complex is trialkylaluminum complex. The ion-exchange activator is (PhNMe<sub>2</sub>H) (B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>).

ABEX EXAMPLE - (2,4,6-Me)<sub>2</sub>N,N-dimethyl-4-aminoazobenzene(Me)EtPh (0.50 g) and (ethylene glycol dimethyl ether)NiBr<sub>2</sub> (0.36 g) were dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (8 ml) under nitrogen and stirred at room temperature for 8 hours. The obtained solution was concentrated and the residue recrystallized from CH<sub>2</sub>Cl<sub>2</sub>/hexane to give (2,4,6-Me)<sub>2</sub>N,N-dimethyl-4-aminoazobenzene(Me)EtPh nickel (II) dibromide (53%).

AN.S DCR-776360

SDCN RABNX7

CM 1

Br




10/551,292

CM 2

Ni

CM 3



AN.S DCR-776360  
SDCN RABNX7


CM 1

Br

CM 2

Ni

CM 3



=> fil hcap wpix

FILE 'HCAPLUS' ENTERED AT 13:41:34 ON 25 JUN 2009  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIX' ENTERED AT 13:41:34 ON 25 JUN 2009  
 COPYRIGHT (C) 2009 THOMSON REUTERS

=> s 145 not 147  
 L48                    5 L45 NOT L47

=> file stnguide

FILE 'STNGUIDE' ENTERED AT 13:41:51 ON 25 JUN 2009  
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
 COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.  
 LAST RELOADED: Jun 19, 2009 (20090619/UP).

=> d ibib ed abs hitind hitstr 1-5  
 YOU HAVE REQUESTED DATA FROM FILE 'HCAPLUS' - CONTINUE? (Y)/N:y

L48 ANSWER 1 OF 5 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2008:732443 HCAPLUS Full-text  
 DOCUMENT NUMBER: 149:257825  
 TITLE: Synthesis of new palladium(II) complexes containing  
 tetradentate-nitrogen donor ligands: Combined  
 structural studies by NMR spectroscopy and X-ray  
 crystallography  
 AUTHOR(S): Espinal, Monica; Pons, Josefina; Garcia-Anton, Jordi;  
 Solans, Xavier; Font-Bardia, Merce; Ros, Josep  
 CORPORATE SOURCE: Departament de Quimica, Unitat de Quimica Inorganica,  
 Universitat Autonoma de Barcelona, Bellaterra,  
 Barcelona, 08193, Spain  
 SOURCE: Inorganica Chimica Acta (2008), 361(9-10), 2648-2658  
 CODEN: ICHAA3; ISSN: 0020-1693  
 PUBLISHER: Elsevier B.V.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 149:257825  
 ED Entered STN: 19 Jun 2008  
 AB Treatment of the tetradentate (NN'N'N) N-alkylaminopyrazole ligands 3,6-  
 dimethyl-1,8-(3,5-dimethyl-1-pyrazolyl)-3,6-diazaoctane (ddad) and 1,4-bis[2-  
 (3,5-dimethyl-1-pyrazolyl)ethyl]piperazine (bedp) with [PdCl<sub>2</sub>(MeCN)<sub>2</sub>] in a 1:1  
 M/L ratio in MeCN produces [PdCl<sub>2</sub>(L)] and [PdCl<sub>2</sub>(L)] (L = ddad and bedp).  
 Treatment of the corresponding complex [PdCl<sub>2</sub>(L)] (L = ddad, bedp) in the  
 presence of AgBF<sub>4</sub> in CH<sub>2</sub>Cl<sub>2</sub>/MeOH (2:1) or NaBF<sub>4</sub> in MeCN gives [Pd(L)](BF<sub>4</sub>)<sub>2</sub>.  
 The Pd(II) complexes were characterized by elemental analyses, conductivity  
 measurements, IR and <sup>1</sup>H and <sup>13</sup>C{<sup>1</sup>H} NMR spectroscopies when possible. The x-  
 ray structure of [Pd(ddad)]Cl<sub>2</sub>·3H<sub>2</sub>O was determined The Pd(II) is coordinated

to the ddad ligand by two N atoms of pyrazolyl groups and two N atoms of the amine groups, in a slightly distorted square-planar geometry.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 75

IT 139775-87-4 139775-88-5


RL: RCT (Reactant); RACT (Reactant or reagent)  
(complexation with palladium)

IT 139775-87-4

RL: RCT (Reactant); RACT (Reactant or reagent)  
(complexation with palladium)

RN 139775-87-4 HCAPLUS

CN 1,2-Ethanediamine, N1,N2-bis[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-N1,N2-dimethyl- (CA INDEX NAME)



REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L48 ANSWER 2 OF 5 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:261803 HCAPLUS Full-text

DOCUMENT NUMBER: 150:144351

TITLE: Synthesis of 1-(2-aminoethyl)pyrazoles under phase-transfer catalysis

AUTHOR(S): Attaryan, O. S.; Baltayan, A. O.; Sagatelyan, R. E.; Takmazyan, K. Ts.

CORPORATE SOURCE: Institute of Organic Chemistry, National Academy of Sciences of Armenia, Yerevan, 375091, Armenia

SOURCE: Russian Journal of General Chemistry (2008), 78(1), 136-138

CODEN: RJGCEK; ISSN: 1070-3632

PUBLISHER: Pleiades Publishing, Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 03 Mar 2008

AB Alkylation of pyrazoles with 2-chloroethylamine was performed under conditions of phase-transfer catalysis. Depending on the substrate acidity, the electrophilic substitution process may be accompanied by dehydrochlorination of the alkylating agent, so that 6 equiv of 2-chloroethylamine should be used in the alkylation of 3,5-dimethylpyrazole.

CC 28-8 (Heterocyclic Compounds (More Than One Hetero Atom))

IT 62821-88-9P 62821-90-3P 101395-71-5P, 1H-Pyrazole-1-ethanamine

101395-72-6P 511513-23-8P 1101099-34-6P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of pyrazoleethanamine by alkylation of pyrazole with chloroethanamine under phase-transfer catalysis)


IT 511513-23-8P 1101099-34-6P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of pyrazoleethanamine by alkylation of pyrazole with chloroethanamine under phase-transfer catalysis)


RN 511513-23-8 HCAPLUS

CN 1,2-Ethanediamine, N1-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]- (CA INDEX

NAME)



RN 1101099-34-6 HCAPLUS

CN 1,2-Ethanediamine, N1-[2-(3-methyl-1H-pyrazol-1-yl)ethyl]- (CA INDEX  
NAME)REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L48 ANSWER 3 OF 5 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:612137 HCAPLUS Full-text

DOCUMENT NUMBER: 146:286880

TITLE: Experimental and theoretical study of lanthanide  
complexes based on linear and macrocyclic  
polyaminopolycarboxylic acids with pyrazolylethyl armsAUTHOR(S): Perez-Mayoral, Elena; Soriano, Elena; Cerdan,  
Sebastian; Ballesteros, PalomaCORPORATE SOURCE: Laboratorio de Sintesis Organica e Imagen Molecular  
por Resonancia Magnetica, Instituto Universitario de  
Investigacion, Facultad de Ciencias, UNED, Madrid,  
E-28040, SpainSOURCE: Proceedings of ECSOC-9, International Electronic  
Conference on Synthetic Organic Chemistry, 9th, Nov.  
1-30, 2005 (2005), C003/1-C003/11. Editor(s): Seijas,  
Julio A.; Vazquez Tato, M. Pilar. Molecular Diversity  
Preservation International: Basel, Switz.  
CODEN: 69IFGU; ISBN: 3-906980-16-2



DOCUMENT TYPE: Conference; (computer optical disk)

LANGUAGE: English


OTHER SOURCE(S): CASREACT 146:286880

ED Entered STN: 26 Jun 2006

GI



- AB The preparation and magnetic relaxometric properties of gadolinium complexes with polyaminopolycarboxylic acids with pyrazolyylethyl arms (I, II) is described. Optimized geometries for the Gd(L)(H<sub>2</sub>O) complexes (L = I, II, diethylenetriamine pentaacetic acid (TPA), and 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid) (DOTA) were calculated from DFT methods.
- CC 78-7 (Inorganic Chemicals and Reactions)  
Section cross-reference(s): 28, 65, 77
- IT 96-32-2 67000-35-5 117499-16-8 149353-23-1 885689-63-4  
927431-26-3  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of polyaminopolycarboxylic acid with pyrazolyylethyl arms)
- IT 885678-69-3P 885689-50-9P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and complexation with gadolinium(III))
- IT 885689-63-4  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of polyaminopolycarboxylic acid with pyrazolyylethyl arms)
- RN 885689-63-4 HCAPLUS
- CN 10-Oxa-2,5,8-triazadodecanoic acid,  
5-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-11,11-dimethyl-9-oxo-,  
1,1-dimethylethyl ester (CA INDEX NAME)





- IT 885689-50-9P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT

(Reactant or reagent)

(preparation and complexation with gadolinium(III))

RN 885689-50-9 HCAPLUS

CN Glycine, N,N'-[[[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]imino]di-2,1-ethanediyl]bis[N-(carboxymethyl)-, sodium salt (1:4) (CA INDEX NAME)



REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L48 ANSWER 4 OF 5 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:601860 HCAPLUS Full-text

DOCUMENT NUMBER: 146:219353

TITLE: Experimental and theoretical study of lanthanide complexes based on linear and macrocyclic polyaminopolycarboxylic acids containing pyrazolyethyl arms

AUTHOR(S): Perez-Mayoral, Elena; Soriano, Elena; Cerdan, Sebastian; Ballesteros, Paloma

CORPORATE SOURCE: Laboratorio de Sintesis Organica e Imagen Molecular por Resonancia Magnetica, Instituto Universitario de Investigacion, Facultad de Ciencias, UNED, Madrid, E-28040, Spain

SOURCE: Molecules (2006), 11(5), 345-356  
CODEN: MOLEFW; ISSN: 1420-3049  
URL: <http://mdpi.org/subscribers/molecules/papers/11050345.pdf>

PUBLISHER: Molecular Diversity Preservation International

DOCUMENT TYPE: Journal; (online computer file)

LANGUAGE: English

OTHER SOURCE(S): CASREACT 146:219353

ED Entered STN: 22 Jun 2006

GI


\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB We report the synthesis of two novel Gd(III)-complexes Gd(L) and Gd(L') derived from linear and macrocyclic polyaminopolycarboxylic acids (I) = H4L and (II) = H3L', resp., and a study of their relaxivity properties. Optimized geometries of the gadolinium complexes also are reported. The relationships between the exptl. and theor. results have provided interesting information about the kinetic and thermodyn. stability of these complexes.


CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 28, 65, 67, 77

- IT 885689-63-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and reactant for preparation of  
 N,N'-[[[(dimethyl-pyrazolyl)ethyl]imino]diethanediyl]bis[N-(carboxymethyl)]glycine)
- IT 885689-50-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and reactant for preparation of gadolinium  
 N,N'-[[[(dimethyl-pyrazolyl)ethyl]imino]diethanediyl]bis[N-(carboxymethyl)]glycine complex)
- IT 885689-67-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (reactant for preparation of N,N'-[[[(dimethyl-pyrazolyl)ethyl]imino]diethanediyl]bis[N-(carboxymethyl)]glycine)
- IT 885689-63-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and reactant for preparation of  
 N,N'-[[[(dimethyl-pyrazolyl)ethyl]imino]diethanediyl]bis[N-(carboxymethyl)]glycine)
- RN 885689-63-4 HCAPLUS
- CN 10-Oxa-2,5,8-triazadodecanoic acid,  
 5-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-11,11-dimethyl-9-oxo-,  
 1,1-dimethylethyl ester (CA INDEX NAME)




- IT 885689-50-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and reactant for preparation of gadolinium  
 N,N'-[[[(dimethyl-pyrazolyl)ethyl]imino]diethanediyl]bis[N-(carboxymethyl)]glycine complex)
- RN 885689-50-9 HCAPLUS
- CN Glycine, N,N'-[[[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]imino]di-2,1-ethanediyl]bis[N-(carboxymethyl)-, sodium salt (1:4) (CA INDEX NAME)



●4 Na

IT 885689-67-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (reactant for preparation of N,N'-[[[(dimethyl-  
 pyrazolyl)ethyl]imino]diethanediyl]bis[N-(carboxymethyl)]glycine)  
 RN 885689-67-8 HCAPLUS  
 CN 2-Oxa-5,8,11-triazatridecan-13-oic acid,  
 8-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-5,11-bis(2-methoxy-2-oxoethyl)-3-  
 oxo-, methyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L48 ANSWER 5 OF 5 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:469891 HCAPLUS Full-text

DOCUMENT NUMBER: 144:459915

TITLE: Preparation of  
 (pyrazolylethyl)diethylenetriaminetetraacetate  
 heterocyclic ligands and their gadolinium(III)  
 complexes with biomedical applications

INVENTOR(S): Ballesteros Garcia, Paloma

PATENT ASSIGNEE(S): Universidad Nacional De Educacion A Distancia  
 (U.N.E.D.), Spain

SOURCE: PCT Int. Appl., 23 pp.  
 CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Spanish

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006051143	A1	20060518	WO 2005-ES602	20051107



W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

ES 2253114 A1 20060516 ES 2004-2679 20041108


ES 2253114 B1 20070701

PRIORITY APPLN. INFO.: ES 2004-2679 A 20041108

OTHER SOURCE(S): MARPAT 144:459915

ED Entered STN: 19 May 2006

GI



AB The invention relates to compds. DTPA-like ligands (I) and their paramagnetic complexes of Gd(III) and other lanthanides which may be used as contrast agents for magnetic resonance imaging. More specifically, the invention relates to compds. I, wherein R1 and/or R2 are hydrogens, Me groups, nitro groups and amino groups. The invention further relates to a method of obtaining said compds. from the corresponding bromoethylpyrazoles, comprising the following steps: 1) alkylation of the original amine; 2) deprotection of the tert-butoxycarbonylamino groups; 3) alkylation of the amino groups with Me bromoacetate; and, finally, 4) basic hydrolysis which produces the tetrasodium salt. The invention relates to complexes of Gd(III) and of other lanthanides derived from compds. I, to the method of obtaining said complexes and to the exptl. and clin. use of same in the production of contrast agents for clin. diagnosis by magnetic resonance imaging. Thus, ligands I (Na4L; X = Na; R1, R2 = H, H; Me, Me; NO2, H; NH2, H) were prepared and complexed with gadolinium to give Na[GdL] complexes. The relaxivities of these gadolinium were measured to determine their suitability as MRI contrast agents.

CC 78-7 (Inorganic Chemicals and Reactions)


Section cross-reference(s): 8, 28

IT 885689-48-5P 885689-50-9P 885689-52-1P 885689-54-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(ligand; preparation of (pyrazolylethyl)diethylenetriaminetetraacetate heterocyclic ligands and their gadolinium(III) complexes for use as MRI

- contrast agents)
- IT 885689-57-6P 885689-59-8P 885689-63-4P  
885689-65-6P 885689-67-8P 885689-69-0P 885689-71-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preparation of (pyrazolylethyl)diethylenetriaminetetraacetate heterocyclic  
 ligands and their gadolinium(III) complexes for use as MRI contrast  
 agents)
- IT 885689-48-5P 885689-50-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (ligand; preparation of (pyrazolylethyl)diethylenetriaminetetraacetate  
 heterocyclic ligands and their gadolinium(III) complexes for use as MRI  
 contrast agents)
- RN 885689-48-5 HCAPLUS
- CN Glycine, N,N'-[[[2-(1H-pyrazol-1-yl)ethyl]imino]di-2,1-ethanediyl]bis[N-(  
 (carboxymethyl)-, tetrasodium salt (9CI) (CA INDEX NAME)



●4 Na

- RN 885689-50-9 HCAPLUS
- CN Glycine, N,N'-[[[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]imino]di-2,1-  
 ethanediyl]bis[N-(carboxymethyl)-, sodium salt (1:4) (CA INDEX NAME)




●4 Na

- IT 885689-57-6P 885689-59-8P 885689-63-4P  
885689-67-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preparation of (pyrazolylethyl)diethylenetriaminetetraacetate heterocyclic  
 ligands and their gadolinium(III) complexes for use as MRI contrast  
 agents)
- RN 885689-57-6 HCAPLUS
- CN 10-Oxa-2,5,8-triazadodecanoic acid,  
 11,11-dimethyl-9-oxo-5-[2-(1H-pyrazol-1-yl)ethyl]-, 1,1-dimethylethyl


10/551,292

ester (CA INDEX NAME)




RN 885689-59-8 HCAPLUS

CN 2-Oxa-5,8,11-triazatridecan-13-oic acid,  
5,11-bis(2-methoxy-2-oxoethyl)-3-oxo-8-[2-(1H-pyrazol-1-yl)ethyl]-, methyl  
ester (9CI) (CA INDEX NAME)



RN 885689-63-4 HCAPLUS


CN 10-Oxa-2,5,8-triazadodecanoic acid,  
5-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-11,11-dimethyl-9-oxo-,  
1,1-dimethylethyl ester (CA INDEX NAME)



RN 885689-67-8 HCAPLUS

CN 2-Oxa-5,8,11-triazatridecan-13-oic acid,  
8-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-5,11-bis(2-methoxy-2-oxoethyl)-3-  
oxo-, methyl ester (9CI) (CA INDEX NAME)

10/551,292



REFERENCE COUNT:

3

THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d que nos 128

```

L5          STR
L7          827 SEA FILE=REGISTRY SSS FUL L5
L10         STR
L12         90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10
L14         QUE SPE=ON ABB=ON PLU=ON SANTOS, I?/AU,AUTH
L15         QUE SPE=ON ABB=ON PLU=ON REGO, I?/AU,AUTH
L16         QUE SPE=ON ABB=ON PLU=ON CORREIA, J?/AU,AUTH
L17         QUE SPE=ON ABB=ON PLU=ON PAULO, A?/AU,AUTH
L18         QUE SPE=ON ABB=ON PLU=ON ALVES, S?/AU,AUTH
L19         QUE SPE=ON ABB=ON PLU=ON VITOR, R?/AU,AUTH
L20         QUE SPE=ON ABB=ON PLU=ON MALLINCKRODT/CS,SO,PA
L22         QUE SPE=ON ABB=ON PLU=ON BIOMOLECUL? OR (BIO(1W)MOLEC
          UL?) OR (BIOLOGIC?(3A)MOLECUL?)
L23         QUE SPE=ON ABB=ON PLU=ON CHELAT?
L24         QUE SPE=ON ABB=ON PLU=ON "CHELATING AGENTS"+PFT,OLD,N
          EW,NT/CT
L25         46 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L12
L26         19 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L25 AND (L22 OR L23
          OR L24)
L27         46 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON (L25 OR L26)
L28         18 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L27 AND (L14 OR L15
          OR L16 OR L17 OR L18 OR L19 OR L20)

```

=> d que nos 137

```

L10         STR
L14         QUE SPE=ON ABB=ON PLU=ON SANTOS, I?/AU,AUTH
L15         QUE SPE=ON ABB=ON PLU=ON REGO, I?/AU,AUTH
L16         QUE SPE=ON ABB=ON PLU=ON CORREIA, J?/AU,AUTH
L17         QUE SPE=ON ABB=ON PLU=ON PAULO, A?/AU,AUTH
L18         QUE SPE=ON ABB=ON PLU=ON ALVES, S?/AU,AUTH
L19         QUE SPE=ON ABB=ON PLU=ON VITOR, R?/AU,AUTH
L20         QUE SPE=ON ABB=ON PLU=ON MALLINCKRODT/CS,SO,PA
L22         QUE SPE=ON ABB=ON PLU=ON BIOMOLECUL? OR (BIO(1W)MOLEC
          UL?) OR (BIOLOGIC?(3A)MOLECUL?)
L23         QUE SPE=ON ABB=ON PLU=ON CHELAT?
L33         10 SEA FILE=WPIX SSS FUL L10
L34         3 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON (RABNX7/DCN OR RAFVJB/DCN
          OR RAFVJC/DCN OR RAFVJD/DCN OR RAFVJE/DCN OR RAFVJF/DCN OR
          RAFVJG/DCN OR RAFVJ8/DCN OR RAFVJ9/DCN OR RAMT8E/DCN) OR
          L33/DCR
L35         2 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON L34 AND (L22 OR L23)
L36         3 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON (L34 OR L35)
L37         1 SEA FILE=WPIX SPE=ON ABB=ON PLU=ON L36 AND (L14 OR L15 OR
          L16 OR L17 OR L18 OR L19 OR L20)

```

=> d que nos 139

```

L5          STR
L7          827 SEA FILE=REGISTRY SSS FUL L5
L10         STR
L12         90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10
L39         0 SEA FILE=MEDLINE SPE=ON ABB=ON PLU=ON L12

```

=> d que nos 141

```

L5          STR
L7          827 SEA FILE=REGISTRY SSS FUL L5

```

```

L10          STR
L12          90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10
L41          0 SEA FILE=EMBASE SPE=ON  ABB=ON  PLU=ON  L12

```

=> d his 142

```

          (FILE 'BIOSIS, CABA, AGRICOLA, BIOTECHNO, DRUGU, VETU' ENTERED AT
          13:30:22 ON 25 JUN 2009)
L42          0 S L12

```

=> d que nos 142

```

L5          STR
L7          827 SEA FILE=REGISTRY SSS FUL L5
L10         STR
L12         90 SEA FILE=REGISTRY SUB=L7 SSS FUL L10
L42         0 SEA L12

```

=> d his 144

```

          (FILE 'MEDLINE, BIOSIS, EMBASE, PASCAL, JAPIO, CABA, CEABA-VTB, LIFESCI,
          BIOENG, BIOTECHNO, BIOTECHDS, DRUGU, DRUGB, VETU, VETB, SCISEARCH,
          CONFSCI, DISSABS, RDISCLOSURE' ENTERED AT 13:31:47 ON 25 JUN 2009)
L44         20 S L43 AND L14-L20

```

=> d que 144

```

L14         QUE  SPE=ON  ABB=ON  PLU=ON  SANTOS, I?/AU,AUTH
L15         QUE  SPE=ON  ABB=ON  PLU=ON  REGO, I?/AU,AUTH
L16         QUE  SPE=ON  ABB=ON  PLU=ON  CORREIA, J?/AU,AUTH
L17         QUE  SPE=ON  ABB=ON  PLU=ON  PAULO, A?/AU,AUTH
L18         QUE  SPE=ON  ABB=ON  PLU=ON  ALVES, S?/AU,AUTH
L19         QUE  SPE=ON  ABB=ON  PLU=ON  VITOR, R?/AU,AUTH
L20         QUE  SPE=ON  ABB=ON  PLU=ON  MALLINCKRODT/CS, SO, PA
L23         QUE  SPE=ON  ABB=ON  PLU=ON  CHELAT?
L43         769 SEA ?PYRAZOL?/IT, TI, CC, CT, ST, STP AND L23/IT, TI, CC, CT, ST, STP
L44         20 SEA L43 AND (L14 OR L15 OR L16 OR L17 OR L18 OR L19 OR L20)

```

=> dup rem 128 137 139 141 142 144

L39 HAS NO ANSWERS

L41 HAS NO ANSWERS

L42 HAS NO ANSWERS

DUPLICATE IS NOT AVAILABLE IN 'RDISCLOSURE'.

ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE

FILE 'HCAPLUS' ENTERED AT 13:43:20 ON 25 JUN 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIX' ENTERED AT 13:43:20 ON 25 JUN 2009

COPYRIGHT (C) 2009 THOMSON REUTERS

FILE 'MEDLINE' ENTERED AT 13:43:20 ON 25 JUN 2009

FILE 'BIOSIS' ENTERED AT 13:43:20 ON 25 JUN 2009

Copyright (c) 2009 The Thomson Corporation

FILE 'EMBASE' ENTERED AT 13:43:20 ON 25 JUN 2009

Copyright (c) 2009 Elsevier B.V. All rights reserved.

FILE 'PASCAL' ENTERED AT 13:43:20 ON 25 JUN 2009  
Any reproduction or dissemination in part or in full,  
by means of any process and on any support whatsoever  
is prohibited without the prior written agreement of INIST-CNRS.  
COPYRIGHT (C) 2009 INIST-CNRS. All rights reserved.

FILE 'BIOENG' ENTERED AT 13:43:20 ON 25 JUN 2009  
COPYRIGHT (C) 2009 Cambridge Scientific Abstracts (CSA)

FILE 'SCISEARCH' ENTERED AT 13:43:20 ON 25 JUN 2009  
Copyright (c) 2009 The Thomson Corporation  
PROCESSING COMPLETED FOR L28  
PROCESSING COMPLETED FOR L37  
PROCESSING COMPLETED FOR L39  
PROCESSING COMPLETED FOR L41  
PROCESSING COMPLETED FOR L42  
PROCESSING COMPLETED FOR L44  
L49           21 DUP REM L28 L37 L39 L41 L42 L44 (18 DUPLICATES REMOVED)  
              ANSWERS '1-18' FROM FILE HCAPLUS  
              ANSWER '19' FROM FILE EMBASE  
              ANSWERS '20-21' FROM FILE SCISEARCH

=> file stnguide

FILE 'STNGUIDE' ENTERED AT 13:43:36 ON 25 JUN 2009  
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.  
LAST RELOADED: Jun 19, 2009 (20090619/UP).

=> d ibib ed abs hitind hitstr 1-18

YOU HAVE REQUESTED DATA FROM FILE 'HCAPLUS, EMBASE, SCISEARCH' - CONTINUE? (Y)/N:y

L49 ANSWER 1 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 1

ACCESSION NUMBER: 2008:1207034 HCAPLUS Full-text

DOCUMENT NUMBER: 150:20350

TITLE: Synthesis, characterization, and evaluation of a novel 99mTc(CO)3 pyrazolyl conjugate of a peptide nucleic acid sequence

AUTHOR(S): Xavier, Catarina; Giannini, Clelia; Gano, Lurdes; Maiorana, Stefano; Alberto, Roger; Santos, Isabel

CORPORATE SOURCE: Unidade de Ciencias Quimicas e Radiofarmaceuticas, Instituto Tecnologico e Nuclear, Sacavem, 2686-953, Port.

SOURCE: JBIC, Journal of Biological Inorganic Chemistry (2008), 13(8), 1335-1344  
CODEN: JJBCFA; ISSN: 0949-8257

PUBLISHER: Springer GmbH

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 08 Oct 2008

AB The 16-mer peptide nucleic acid sequence H-A GAT CAT GCC CGG CAT-Lys-NH<sub>2</sub> (1), which is complementary to the translation start region of the N-myc oncogene mRNA, was synthesized and conjugated to a pyrazolyl diamine bifunctional chelator (pz). The novel conjugate pz-A GAT CAT GCC CGG CAT-Lys-NH<sub>2</sub> (2) was labeled with technetium tricarbonyl, yielding quant. the complex fac-[99mTc(CO)<sub>3</sub>(κ<sup>3</sup>-pz-A GAT CAT GCC CGG CAT-Lys-NH<sub>2</sub>)]<sup>2+</sup> (4). Complex 4 was obtained with high radiochem. purity and high specific activity, revealing high stability in human serum and in cell culture medium. The identity of 4 was confirmed by comparing its reversed-phase high performance liquid chromatog. profile with that of the rhenium analog fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-pz-A GAT CAT GCC CGG CAT-Lys-NH<sub>2</sub>)]<sup>2+</sup> (3), prepared by conjugation of fac-[Re(CO)<sub>3</sub>(3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>N((CH<sub>2</sub>)<sub>3</sub>COOH)(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub>)]<sup>+</sup> to 1, using solid-phase techniques. UV melting expts. of 1 and 3 with the complementary DNA sequence led to the formation of stable duplexes, indicating that the conjugation of 1 to the pyrazolyl chelator and to the metal fragment fac-[M(CO)<sub>3</sub>]<sup>+</sup> did not affect the recognition of the complementary sequence as well as the duplex stability. For a first screening, SH-SY5Y human neuroblastoma cells, which express N-myc, were treated with 4. The results show that 4 internalizes (7% of the activity goes into the cells, after 4 h at 37°), presenting also a relatively high cellular retention (only 40% of internalized activity is released from the cells after 5 h).

CC 34-3 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 8, 78

IT Nucleic acid hybridization

(DNA-PNA; solid-phase preparation of peptide nucleic acid sequences, their conjugates with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

IT Imaging agents

(solid-phase preparation of peptide nucleic acid sequences, their conjugates with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

IT Peptide nucleic acids

RL: BSU (Biological study, unclassified); PRP (Properties); RCT



(Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)  
 (solid-phase preparation of peptide nucleic acid sequences, their conjugates  
 with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

IT 1089234-29-6P  
 RL: BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (solid-phase preparation of peptide nucleic acid sequences, their conjugates  
 with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

IT 1089234-31-0 1089234-32-1  
 RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)  
 (solid-phase preparation of peptide nucleic acid sequences, their conjugates  
 with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

IT 1089234-26-3P 1089234-28-5P  
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)  
 (solid-phase preparation of peptide nucleic acid sequences, their conjugates  
 with pyrazolyl chelator, and complexes with rhenium and 99-technetium)


IT 1089234-27-4P  
 RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (solid-phase preparation of peptide nucleic acid sequences, their conjugates  
 with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

IT 163932-31-8 782501-78-4 945384-90-7  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (solid-phase preparation of peptide nucleic acid sequences, their conjugates  
 with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

IT 1089234-30-9DP, resin-bound  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (solid-phase preparation of peptide nucleic acid sequences, their conjugates  
 with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

IT 782501-78-4  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (solid-phase preparation of peptide nucleic acid sequences, their conjugates  
 with pyrazolyl chelator, and complexes with rhenium and 99-technetium)

RN 782501-78-4 HCAPLUS  
 CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



REFERENCE COUNT: 55 THERE ARE 55 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 2 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 2

ACCESSION NUMBER: 2008:521195 HCAPLUS Full-text

DOCUMENT NUMBER: 150:578080

TITLE: Pyrazolyl-diamine ligands that bear anthracenyl moieties and their rhenium(I) tricarbonyl complexes: synthesis, characterization and DNA-binding properties

AUTHOR(S): Vitor, Rute F.; Correia, Isabel; Videira, Margarida; Marques, Fernanda; Paulo, Antonio; Pessoa, Joao Costa; Viola, Giampietro; Martins, Gabriel G.; Santos, Isabel

CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem, 2686-953, Port.

SOURCE: ChemBioChem (2008), 9(1), 131-142

CODEN: CBCHFX; ISSN: 1439-4227

PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 30 Apr 2008

AB Two novel families of pyrazolyl-diamine ligands that bear an anthracen-9-yl group as a DNA-binding fragment, pz\*(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NHCH<sub>2</sub>-9-anthryl (pz\* = pz (L1), 3,5-Me<sub>2</sub>pz (L2)) and pz\*(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub> (pz\* = 4-(9-anthrylmethyl)pz (L3), 3,5-Me<sub>2</sub>-4-(9-anthrylmethyl)pz (L4)), were prepared and fully characterized. In the case of L2-L4, the evaluation of their coordination capability towards the fac-[Re(CO)<sub>3</sub>]+ core gave the organometallic complexes fac-[Re(CO)<sub>3</sub>{3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NHCH<sub>2</sub>-9-anthryl}]Br (7) and fac-[Re(CO)<sub>3</sub>{4-(9-anthrylmethyl)pz\*(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub>}]Br (pz\* = pz (8), 3,5-Me<sub>2</sub>pz (9)). The interaction of the novel pyrazole-diamine ligands and the rhenium(I) complexes with calf thymus (CT) DNA was studied with a variety of spectroscopic techniques (UV-visible, fluorescence, CD and linear dichroism (LD)). All of the evaluated compds. have a moderate affinity to CT DNA (3.46 + 10<sup>3</sup> < K<sub>b</sub> < 1.95 + 10<sup>4</sup>), but the binding mode depends on the position of the chromophore in the framework of the pyrazolyl-diamine ligands. LD measurements showed that L1 and L2 act as DNA intercalators, but complex 7 intercalates only partially. By contrast, the compds. with the anthracenyl group at the 4-position of the azolyl ring (L3, L4 and 9) do not intercalate, and behave more like DNA groove binders. Fluorescence microscopy studies demonstrated that complexes 7 and 9 can target the nucleus of murine B16-F1 melanoma cells, and appear to be promising platforms for the further design of radiopharmaceuticals for targeted radiotherapy.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 6, 28

IT 944389-67-7P 1152004-05-1P 1152004-07-3P  
1152004-10-8P


RL: BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)

(preparation, complexation with rhenium(I), and DNA binding)


10/551,292

IT 107-15-3, Ethylenediamine, reactions 109-84-2, 2-Hydroxyethylhydrazine  
123-54-6, Acetylacetone, reactions 642-31-9, 9-Anthracenecarboxaldehyde  
2417-77-8, 9-Bromomethylantracene 58353-41-6,  
2-(9-Anthrylmethyl)propane-1,3-diol 511513-23-8  
782501-70-6  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reactant for preparation of substituted pyrazolyldiamine)  
IT 1152004-05-1F 1152004-07-3F  
RL: BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic  
preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant  
or reagent)  
(preparation, complexation with rhenium(I), and DNA binding)  
RN 1152004-05-1 HCAPLUS  
CN 1,2-Ethanediamine, N1-(9-anthracenylmethyl)-N2-[2-(3,5-dimethyl-1H-pyrazol-  
1-yl)ethyl]- (CA INDEX NAME)



PAGE 1-A



PAGE 2-A



RN 1152004-07-3 HCAPLUS  
CN 1,2-Ethanediamine, N1-(9-anthracenylmethyl)-N2-[2-(1H-pyrazol-1-yl)ethyl]-  
(CA INDEX NAME)




IT 511513-23-8 782501-70-6

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant for preparation of substituted pyrazolyldiamine)


RN 511513-23-8 HCAPLUS

CN 1,2-Ethanediamine, N1-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]- (CA INDEX NAME)



RN 782501-70-6 HCAPLUS

CN 1,2-Ethanediamine, N1-[2-(1H-pyrazol-1-yl)ethyl]- (CA INDEX NAME)



REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 3 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 3

ACCESSION NUMBER: 2007:1476534 HCAPLUS Full-text

DOCUMENT NUMBER: 149:241344

TITLE: A 99mTc(CO)3-labeled

pyrazolyl- $\alpha$ -melanocyte-stimulating hormone  
analog conjugate for melanoma targeting

AUTHOR(S): Raposinho, Paula D.; Correia, Joao D. G.;  
Alves, Susana; Botelho, Maria F.; Santos, Ana  
C.; Santos, Isabel

CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem, 2686-953, Port.

SOURCE: Nuclear Medicine and Biology (2008), 35(1), 91-99

CODEN: NMBIEO; ISSN: 0969-8051

PUBLISHER: Elsevier Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 28 Dec 2007

AB Melanoma primary tumors can be, in most cases, removed surgically, whereas there is no satisfactory treatment for metastatic melanoma, being almost always lethal at this stage. Therefore, early detection of primary melanoma tumors is essential. The finding that melanocortin-1 receptor (MC1R) is overexpressed in isolated melanoma cells and melanoma tissues led to the radiolabeling of several  $\alpha$ -MSH ( $\alpha$ -MSH) analogs for early detection and treatment of melanoma. We have coupled the  $\alpha$ -MSH analog Ac-Nle-Asp-His-D-Phe-Arg-Trp-Gly-Lys-NH<sub>2</sub>, through the  $\epsilon$ -amino group of Lys11, to a pyrazolyl-containing chelator (pz). The resulting pz- $\alpha$ -MSH analog reacted with the fac-[99mTc(CO)3]+ moiety, giving [Ac-Nle<sup>4</sup>,Asp<sup>5</sup>,D-Phe<sup>7</sup>,Lys<sup>11</sup>(pz- 99mTc(CO)3)] $\alpha$ -MSH<sub>4-11</sub> in high yield, high specific activity and high radiochem. purity. This radioconjugate, which presents remarkable stability in vitro, exhibited time- and temperature-dependent internalization (4 h at 37°C; 56.7% maximum internalization) and high cellular retention (only 38% was released from the cell after 5 h) in murine melanoma B16F1 cells. A significant tumor uptake [4.2  $\pm$  0.9%ID/g, at 4 h postinjection (p.i.)] was also obtained in melanoma-bearing C57BL6 mice. The in vivo affinity and specificity of the radioconjugate to MC1R were demonstrated by receptor-blocking studies with the potent NDP-MSH agonist (63.5% reduction in tumor uptake at 4 h p.i.).

CC 8-9 (Radiation Biochemistry)

IT 163932-31-8 1044531-54-5

RL: RCT (Reactant); RACT (Reactant or reagent)

(99mTc(CO)3-labeled pyrazolyl- $\alpha$ -MSH analog conjugate for melanoma  
imaging)

IT 1044531-54-5



RL: RCT (Reactant); RACT (Reactant or reagent)

(99mTc(CO)3-labeled pyrazolyl- $\alpha$ -MSH analog conjugate for melanoma  
imaging)

RN 1044531-54-5 HCAPLUS

CN L-Lysinamide, N-acetyl-L-norleucyl-L- $\alpha$ -aspartyl-L-histidyl-D-phenylalanyl-L-arginyl-L-tryptophylglycyl-N<sub>6</sub>-[3-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]propyl]- (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 4 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 4  
 ACCESSION NUMBER: 2007:159588 HCAPLUS Full-text  
 DOCUMENT NUMBER: 146:413668  
 TITLE: Evaluation of two chelators for labeling a  
 PNA monomer with the fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> moiety  
 AUTHOR(S): Xavier, Catarina; Pak, Jae-Kyoung; Santos,  
Isabel; Alberto, Roger  
 CORPORATE SOURCE: Departamento de Quimica, Instituto Tecnológico e  
 Nuclear, Estrada Nacional 10, Sacavem, 2686-953, Port.  
 SOURCE: Journal of Organometallic Chemistry (2007), 692(6),  
 1332-1339  
 CODEN: JORCAI; ISSN: 0022-328X  
 PUBLISHER: Elsevier Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 146:413668  
 ED Entered STN: 13 Feb 2007  
 GI

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*


- AB A PNA monomer containing thymine as nucleobase was synthesized, characterized and coupled to the pyrazolyl containing ligand 3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>N((CH<sub>2</sub>)<sub>3</sub>COOH)(CH<sub>2</sub>)<sub>2</sub>NHBoc and to a modified cysteine S-(carboxymethyl-pentafluorophenyl)-N-[(trifluoro)carbonyl]-cysteine Me ester yielding the bifunctional chelators I and II, resp. Reactions of I and II with the Re(I) tricarbonyl starting material [Re(CO)<sub>3</sub>(H<sub>2</sub>O)<sub>3</sub>]Br afforded the complexes fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-I)]<sup>+</sup> and fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-II)], resp. The identities of the rhenium complexes have been established based on IR spectroscopy, elemental anal., ESI-MS spectrometry and HPLC. The multinuclear NMR spectroscopy (1H, 13C, g-COSY, g-HSQC) has also been very informative in the case of complex fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-I)]<sup>+</sup>, showing the presence of rotamers in solution For fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-II)] the NMR spectrum was too complex due to the presence of rotamers and diastereoisomers. The radioactive congeners of the rhenium complexes, fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>(κ<sup>3</sup>-6)]<sup>+</sup> and fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>(κ<sup>3</sup>-7)], have been prepared by reacting the precursor fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>(H<sub>2</sub>O)<sub>3</sub>]<sup>+</sup> with the corresponding ligands and their identities were established by comparing their HPLC chromatograms with those of the rhenium analogs.
- CC 78-7 (Inorganic Chemicals and Reactions)  
Section cross-reference(s): 26
- ST pyrazolyl deriv thymynyl glycinate chelator prepn complexation rhenium technetium; cysteinyl deriv thymynyl glycinate chelator prepn complexation rhenium technetium; rhenium thymynyl glycinate pyrazolyl cysteinyl deriv carbonyl complex prepn; technetium thymynyl glycinate pyrazolyl cysteinyl deriv carbonyl complex prepn; peptide nucleic acid monomer thymine deriv prepn coupling
- IT Transition metal complexes  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(cysteine-containing peptide; preparation of pyrazolyl-/cysteinyl-containing thymynyl glycinate derivative bifunctional chelators and their rhenium/technetium carbonyl complexes)
- IT Peptides, preparation  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(cysteine-containing, transition metal complexes; preparation of pyrazolyl-/cysteinyl-containing thymynyl glycinate derivative bifunctional chelators and their rhenium/technetium carbonyl complexes)
- IT Transition metal complexes  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(peptide; preparation of pyrazolyl-/cysteinyl-containing thymynyl glycinate derivative bifunctional chelators and their rhenium/technetium carbonyl complexes)
- IT Peptide nucleic acids  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of pyrazolyl-/cysteinyl-containing thymynyl glycinate derivative bifunctional chelators and their rhenium/technetium carbonyl complexes)
- IT Peptides, preparation  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(transition metal complexes; preparation of pyrazolyl-/cysteinyl-containing thymynyl glycinate derivative bifunctional chelators and their rhenium/technetium carbonyl complexes)
- IT 5292-43-3, tert-Butylbromoacetate 14533-84-7 20924-05-4,  
Thymin-1-ylacetic acid 55757-46-5, N-(tert-Butoxycarbonyl)-L-cysteine methyl ester 128421-86-3 163932-31-8 782501-78-4  
828915-71-5

RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation of pyrazolyl-/cysteinyl-containing thyminylyl glycinate derivative  
 bifunctional chelators and their rhenium/technetium carbonyl complexes)  
 IT 24997-00-0P 152774-08-8P 933789-23-2P 933789-24-3P  
933789-25-4P 933789-27-6P 933789-33-4P 1005462-11-2P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preparation of pyrazolyl-/cysteinyl-containing thyminylyl glycinate derivative  
 bifunctional chelators and their rhenium/technetium carbonyl complexes)  
 IT 933789-29-8P 933789-32-3P 933789-34-5P 933789-35-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of pyrazolyl-/cysteinyl-containing thyminylyl glycinate derivative  
 bifunctional chelators and their rhenium/technetium carbonyl complexes)  
 IT 782501-78-4  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation of pyrazolyl-/cysteinyl-containing thyminylyl glycinate derivative  
 bifunctional chelators and their rhenium/technetium carbonyl complexes)  
 RN 782501-78-4 HCAPLUS  
 CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



IT 933789-25-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preparation of pyrazolyl-/cysteinyl-containing thyminylyl glycinate derivative  
 bifunctional chelators and their rhenium/technetium carbonyl complexes)  
 RN 933789-25-4 HCAPLUS  
 CN Glycine, N-[2-[[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]amino]ethyl]-N-[2-(3,4-dihydro-5-methyl-2,4-dioxo-1(2H)-pyrimidinyl)acetyl]- (CA INDEX NAME)





— Me

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 5 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 5

ACCESSION NUMBER: 2008:220414 HCAPLUS Full-text

DOCUMENT NUMBER: 150:121723

TITLE: A pyrazolylamine-phosphonate monoester chelator for the fac-[M(CO)<sub>3</sub>]<sup>+</sup> core (M = Re, <sup>99m</sup>Tc): synthesis, coordination properties and biological assessment

AUTHOR(S): Palma, Elisa; Oliveira, Bruno L.; Figueira, Flavio; Correia, Joao D. G.; Raposinho, Paula D.; Santos, Isabel

CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem, 2686-953, Port.  
SOURCE: Journal of Labelled Compounds and Radiopharmaceuticals (2007), 50(13), 1176-1184

CODEN: JLCRD4; ISSN: 0362-4803


PUBLISHER: John Wiley & Sons Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 21 Feb 2008

GI



AB Rhenium and technetium tricarbonyl pyrazolyl phosphonates I (6, 6a; M = Re, <sup>99m</sup>Tc; X = Et) were prepared as radioimaging or radiotherapeutic agents and probed for biodistribution and biostability in a number of mice organs. Aiming to develop new strategies for the labeling of hydroxyl-containing biomols. with the organometallic core fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup>, a new model bifunctional chelator, Et hydrogen (2-{[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino}ethyl)phosphonate (L4), combining a pyrazolyl-amine chelating

group and a monophosphonate Et ester function ( $-P(O)OHOEt$ ). The phosphonate group allows metal stabilization, and, simultaneously, can be considered as a potential attachment site for a biomol. Reaction of L4 with the precursor  $[^{99m}Tc(H_2O)_3(CO)_3]^+$  gave the model radiocomplex I,  $[^{99m}Tc(CO)_3(k^3-L4)]$  (6a). This radiocomplex was identified by comparing its chromatog. profile with that of the corresponding Re analog 6. Radiocomplex 6a is moderately lipophilic ( $\log P_{o/w} = 1.07$ ), presenting high stability in vitro without any measurable decomposition or ligand exchange, even in the presence of strong competing chelators such as histidine and cysteine at  $37^\circ$  for 24 h. Biodistribution studies of the complex in CD-1 mice indicated a rapid blood clearance, and a rapid clearance from main organs, occurring primarily through the hepatobiliary pathway. Complex 6a presents also a high robustness in vivo, demonstrated by its resistance to metabolic degradation in blood, and intact excretion into the urine, after RP-HPLC anal. of blood and urine samples. Hydrolyzed forms of I ( $X = H$ ) can be coupled with hydroxyl-containing biomols. as a phosphonate ester, thus allowing radioactive labeling.

CC 29-7 (Organometallic and Organometalloidal Compounds)

Section cross-reference(s): 63, 78

ST technetium rhenium chelate pyrazolyl aminophosphonate prepn  
biodistribution biostability; radioactive labeling agent prepn technetium  
chelate tricarbonyl pyrazolyl aminophosphonate; pharmacokinetics  
radioactive labeling agent technetium chelate tricarbonyl  
pyrazolyl aminophosphonate

IT Chelates

RL: PKT (Pharmacokinetics); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(conjugates, technetium, rhenium; preparation, biodistribution and biostability of rhenium and technetium-99m tricarbonyl pyrazolyl aminophosphonate chelates as radioimaging and radiotherapy agents)

IT Partition

(octanol-water; preparation, biodistribution and biostability of rhenium

and

technetium-99m tricarbonyl pyrazolyl aminophosphonate chelates  
as radioimaging and radiotherapy agents)

IT Bioavailability

Chelating agents

Isotope indicators

Lipophilicity

Pharmacokinetics

Radiography

Radiotherapy

(preparation, biodistribution and biostability of rhenium and technetium-

99m

tricarbonyl pyrazolyl aminophosphonate chelates as  
radioimaging and radiotherapy agents)

IT Group VIIIB element complexes

RL: PKT (Pharmacokinetics); RCT (Reactant); SPN (Synthetic preparation);

BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)

(preparation, biodistribution and biostability of rhenium and technetium-

99m

tricarbonyl pyrazolyl aminophosphonate chelates as  
radioimaging and radiotherapy agents)


IT Chelates

Phosphonates


RL: PKT (Pharmacokinetics); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(technetium, rhenium; preparation, biodistribution and biostability of rhenium and technetium-99m tricarbonyl pyrazolyl aminophosphonate chelates as radioimaging and radiotherapy agents)


IT 5324-30-1, Phosphonic acid, 2-bromoethyl-, diethyl ester 67000-35-5  
 524744-56-7 828915-71-5, Rhenium(1+), triaquatricarbonyl-, bromide  
 (OC-6-22)-  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation, biodistribution and biostability of rhenium and technetium-  
 99m  
 tricarbonyl pyrazolyl aminophosphonate chelates as  
 radioimaging and radiotherapy agents)  
 IT 62821-88-9P 144369-80-2P 1096702-45-2P 1096702-46-3P  
1096702-50-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preparation, biodistribution and biostability of rhenium and technetium-  
 99m  
 tricarbonyl pyrazolyl aminophosphonate chelates as  
 radioimaging and radiotherapy agents)  
 IT 1096702-47-4P 1096702-48-5P 1097641-45-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation, biodistribution and biostability of rhenium and technetium-  
 99m  
 tricarbonyl pyrazolyl aminophosphonate chelates as  
 radioimaging and radiotherapy agents)  
 IT 1096702-49-6P  
 RL: BSU (Biological study, unclassified); PKT (Pharmacokinetics); SPN  
 (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (99mTc complex; preparation, biodistribution and biostability of rhenium  
 and  
 technetium-99m tricarbonyl pyrazolyl aminophosphonate chelates  
 as radioimaging and radiotherapy agents)  
 IT 1096702-45-2P 1096702-46-3P 1096702-50-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preparation, biodistribution and biostability of rhenium and technetium-  
 99m  
 tricarbonyl pyrazolyl aminophosphonate chelates as  
 radioimaging and radiotherapy agents)  
 RN 1096702-45-2 HCAPLUS  
 CN Phosphonic acid, P-[2-[[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]ethyl]-  
 , diethyl ester (CA INDEX NAME)



RN 1096702-46-3 HCAPLUS  
 CN 1H-Pyrazole-1-ethanamine, 3,5-dimethyl-N-(2-phosphinoethyl)- (CA INDEX  
 NAME)



RN 1096702-50-9 HCAPLUS  
 CN Phosphonic acid, P-[2-[[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]ethyl]-, monoethyl ester (CA INDEX NAME)



REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 6 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 6

ACCESSION NUMBER: 2007:22645 HCAPLUS [Full-text](#)

DOCUMENT NUMBER: 146:307704

TITLE: Rhenium(V) oxocomplexes with novel pyrazolyl-based N4- and N3S-donor chelators

AUTHOR(S): Moura, Carolina; Vitor, Rute F.; Maria, Leonor; Paulo, Antonio; Santos, Isabel C.; Santos, Isabel

CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem, 2686-953, Port.

SOURCE: Dalton Transactions (2006), (47), 5630-5640

CODEN: DTARAF; ISSN: 1477-9226

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 146:307704

ED Entered STN: 08 Jan 2007

AB The novel pyrazolyl-based ligands 3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub> (1) and pz\*(CH<sub>2</sub>)<sub>2</sub>NH-Gly-CH<sub>2</sub>St<sub>3</sub> (pz\* = pz (8), 3,5-Me<sub>2</sub>pz (9), 4-(EtOOC)CH<sub>2</sub>-3,5-Me<sub>2</sub>pz (10)) were synthesized, and their suitability to stabilize Re(V) oxocomplexes was evaluated using different starting materials, (NBu<sub>4</sub>)[ReOCl<sub>4</sub>], [ReOCl<sub>3</sub>(PPh<sub>3</sub>)<sub>2</sub>] and trans-[ReO<sub>2</sub>(py)<sub>4</sub>]Cl. Compound 1 reacts with trans-[ReO<sub>2</sub>(py)<sub>4</sub>]Cl yielding the cationic compound [ReO(OMe){3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>N(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub>}](BPh<sub>4</sub>) (11) in a low isolated yield. In contrast, the neutral complexes [ReO{pz\*(CH<sub>2</sub>)<sub>2</sub>NH-Gly-CH<sub>2</sub>S}] (pz\* = pz (12), 3,5-Me<sub>2</sub>pz (13), 4-(EtOOCCH<sub>2</sub>)-3,5-Me<sub>2</sub>pz (14)) were synthesized almost quantitatively by reacting [ReOCl<sub>3</sub>(PPh<sub>3</sub>)<sub>2</sub>] or (NBu<sub>4</sub>)[ReOCl<sub>4</sub>] with the trityl-protected chelators 8-10. The x-ray diffraction anal. of 11 and 13 confirmed the tetradentate coordination mode of the resp. ancillary ligands. In 11 the monoanionic chelator coordinates to the metal through four N atoms, while in 13 the chelator is trianionic, coordinating to the metal through three nitrogens and one S atom. Solution NMR studies of 12-14, including two-dimensional NMR techniques (1H COSY and 1H/13C HSQC), confirmed that the N3S coordination mode of the chelators is retained in solution. Unlike 11, complexes 12-14 may be considered relevant in the development of radiopharmaceuticals, as further

corroborated by the synthesis of the congener [99mTcO{pz(CH<sub>2</sub>)<sub>2</sub>-NH-Gly-CH<sub>2</sub>S}] (12a). This radioactive compound was obtained from 99mTcO<sub>4</sub><sup>-</sup> in aqueous medium, in almost quant. yield and with high specific activity and radiochem. purity.


- CC 78-7 (Inorganic Chemicals and Reactions)  
Section cross-reference(s): 8, 75
- IT Crystal structure  
Molecular structure  
(of oxorhenium pyrazolyl-based N4- and N3S-donor chelating ligand complexes)
- IT 927883-68-9P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(for preparation of pyrazolyl-based N3S-donor chelating ligand and its oxorhenium complex)
- IT 302-01-2, Hydrazine, reactions 1074-82-4, Potassium phthalimide 1972-28-7, Diethylazodicarboxylate 7087-68-5, DIPEA 67000-35-5 91425-33-1 119291-22-4 927883-81-6  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of pyrazolyl-based N3S-donor chelating ligand and its oxorhenium complexes)
- IT 62821-88-9P 101395-71-5P, 1H-Pyrazole-1-ethanamine 121751-71-1P 144369-80-2P 927883-67-8P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(for preparation of pyrazolyl-based N3S-donor chelating ligand and its oxorhenium complexes)
- IT 67000-34-4  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of pyrazolyl-based N4-chelating ligand and its oxorhenium complexes)
- IT 383-63-1P, Ethyl trifluoroacetate  
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
(for preparation of pyrazolyl-based N4-donor chelating ligand and its oxorhenium complexes)
- IT 111-40-0 24424-99-5, Di-tert-butyl dicarbonate 120131-72-8  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of pyrazolyl-based N4-donor chelating ligand and its oxorhenium complexes)
- IT 53675-30-2, Tetrabutylammonium tetrachlorooxorhenate  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of rhenium oxo pyrazolyl-based N3S-donor chelating ligand complex)
- IT 17442-18-1, Trichloro(oxo)bis(triphenylphosphine)rhenium  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of rhenium oxo pyrazolyl-based N3S-donor chelating ligand complexes)
- IT 31429-86-4, trans-[Dioxotetrakis(pyridine)]rhenium(1+) chloride  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of rhenium oxo pyrazolyl-based N4-donor chelating ligand complex)
- IT 23288-60-0, Sodium tetraoxotechnetate(1-)-99Tc  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of technetium-99m pyrazolyl-based N4-donor chelating ligand complex)
- IT 927883-66-7P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and reaction with dioxorhenium pyridine complex)
- IT 927883-66-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and reaction with dioxorhenium pyridine complex)

RN 927883-66-7 HCAPLUS

CN 1,2-Ethanediamine, N1-(2-aminoethyl)-N2-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]- (CA INDEX NAME)



REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 7 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 7

ACCESSION NUMBER: 2006:711846 HCAPLUS Full-text

DOCUMENT NUMBER: 146:223509

TITLE: Pyrazolyl conjugates of bombesin: A new tridentate ligand framework for the stabilization of fac-[M(CO)<sub>3</sub>]<sup>+</sup> moiety

AUTHOR(S): Alves, Susana; Correia, Joao D. G.  
; Santos, Isabel; Veerendra, Bhadrasetty;  
Sieckman, Gary L.; Hoffman, Timothy J.; Rold, Tammy L.; Figueroa, Said Daibes; Retzlöff, Lauren; McCrate, Joseph; Prasanphanich, Adam; Smith, Charles J.

CORPORATE SOURCE: Department of Radiology, University of Missouri-Columbia School of Medicine, Columbia, MO, 65211, USA

SOURCE: Nuclear Medicine and Biology (2006), 33(5), 625-634  
CODEN: NMBIEO; ISSN: 0969-8051

PUBLISHER: Elsevier Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 21 Jul 2006

AB We have described the synthesis of tridentate pyrazolyl ligand frameworks for coordination to the fac-[\*M(CO)<sub>3</sub>]<sup>+</sup> metal fragment (\*M = 186/188Re or 99mTc). These ligands impart a degree of kinetic inertness on the metal center, warranting their study in biol. systems. We herein report in vitro/in vivo radiolabeling investigations of a new series of pyrazolyl bombesin (BBN) conjugates radiolabeled via the Isolink kit. These new conjugates are based on the general structure [99mTc-pyrazolyl-X-BBN[7-14]NH<sub>2</sub>], where X = β-alanine, serylserylserine or glycylglycylglycine. The pyrazolyl ligand is a tridentate ligand framework that coordinates the metal center through nitrogen donor atoms. The results of these investigations demonstrate the ability of these new conjugates to specifically target the gastrin-releasing peptide receptor subtype 2, which is overexpressed on human prostate PC-3 cancerous tissues. Therefore, these studies suggest the tridentate pyrazolyl ligand framework to be an ideal candidate for the design and development of low-valent 99mTc-based diagnostic radiopharmaceuticals based on BBN or other targeting vectors.

CC 8-9 (Radiation Biochemistry)

IT 163932-31-8 924660-90-2 924660-91-3  
924660-92-4

RL: RCT (Reactant); RACT (Reactant or reagent)

(pyrazolyl conjugates of bombesin: new tridentate ligand framework for stabilization of fac-[M(CO)<sub>3</sub>]<sup>+</sup> moiety)

IT 924660-90-2 924660-91-3 924660-92-4


RL: RCT (Reactant); RACT (Reactant or reagent)

(pyrazolyl conjugates of bombesin: new tridentate ligand framework for stabilization of fac-[M(CO)<sub>3</sub>]<sup>+</sup> moiety)

RN 924660-90-2 HCAPLUS


CN L-Methioninamide, N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]glycylglycylglycyl-L-glutamyl-L-tryptophyl-L-alanyl-L-valylglycyl-L-histidyl-L-leucyl- (CA INDEX NAME)

Absolute stereochemistry.



PAGE 1-B

—SMe






RN 924660-91-3 HCAPLUS


CN L-Methioninamide, N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-L-seryl-L-seryl-L-seryl-L-glutamyl-L-tryptophyl-L-alanyl-L-valylglycyl-L-histidyl-L-leucyl- (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B








RN 924660-92-4 HCAPLUS



CN L-Methioninamide, N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-β-alanyl-L-glutaminy-L-tryptophyl-L-alanyl-L-valylglycyl-L-histidyl-L-leucyl- (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



H<sub>2</sub>N

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 8 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 8

ACCESSION NUMBER: 2005:205084 HCAPLUS Full-text

DOCUMENT NUMBER: 142:406796

TITLE: Pyrazolyl Derivatives as Bifunctional  
Chelators for Labeling Tumor-Seeking Peptides  
with the fac-[M(CO)<sub>3</sub>]<sup>+</sup> Moiety (M = <sup>99m</sup>Tc, Re):  
Synthesis, Characterization, and Biological Behavior  
AUTHOR(S): Alves, Susana; Paulo, Antonio;  
Correia, Joao D. G.; Gano, Lurdes; Smith,  
Charles J.; Hoffman, Timothy J.; Santos,  
Isabel

CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem, 2686-953, Port.

SOURCE: Bioconjugate Chemistry (2005), 16(2), 438-449

CODEN: BCCHE\$; ISSN: 1043-1802

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English


ED Entered STN: 09 Mar 2005

AB Radiolabeling of biol. active moles with the [<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> unit has been of primary interest in recent years. With this in mind, we herein report sym. (L1) and asym. (L2-L5) pyrazolyl-containing chelators that have been evaluated in radiochem. reactions with the synthon [<sup>99m</sup>Tc(H<sub>2</sub>O)<sub>3</sub>(CO)<sub>3</sub>]<sup>+</sup> (1a). These reactions yielded the radioactive building blocks [<sup>99m</sup>Tc(CO)<sub>3</sub>(κ<sup>3</sup>-L)]<sup>+</sup> (L = L1-L5, 2a-6a), which were identified by RP-HPLC. The corresponding Re surrogates (2-6) allowed for macroscopic identification of the radiochem. conjugates. Complexes 2a-6a, with log P<sub>o/w</sub> values ranging from -2.35 to 0.87, were obtained in yields of ≥90% using ligand concns. in the 10<sup>-5</sup>-10<sup>-4</sup> M range. Challenge studies with cysteine and histidine revealed high stability for all of these radioactive complexes, and biodistribution studies in mice indicated a fast rate of blood clearance and high rate of total radioactivity excretion, occurring primarily through the renal-urinary pathway. Based on the framework of the asym. chelators, the novel bifunctional ligands 3,5-Me<sub>2</sub>-pz(CH<sub>2</sub>)<sub>2</sub>N((CH<sub>2</sub>)<sub>3</sub>COOH)(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub> (L6) and pz(CH<sub>2</sub>)<sub>2</sub>N((CH<sub>2</sub>)<sub>3</sub>COOH)(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub> (L7) have been synthesized and their coordination chemical toward (NEt<sub>4</sub>)<sub>2</sub>[ReBr<sub>3</sub>(CO)<sub>3</sub>] (1) has been explored. The resulting complexes, fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-L)]Br (L6 (7), L7 (8)), contain tridentate ancillary ligands that are coordinated to the metal center through the pyrazolyl and amine nitrogen atoms, as observed for the other related building blocks. L6 and L7 were coupled to a glycylglycine Et ester dipeptide, and the resulting functionalized ligands were used to prepare the model complexes fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-3,5-Me<sub>2</sub>-pz(CH<sub>2</sub>)<sub>2</sub>N(glygly)(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub>)]<sup>+</sup> (9/9a) and fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-pz(CH<sub>2</sub>)<sub>2</sub>N(CH<sub>2</sub>)<sub>3</sub>(glygly)(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub>)]<sup>+</sup> (10/10a) (M = Re, <sup>99m</sup>Tc). These small conjugates have been fully characterized and are reported herein. On the basis of the in vitro/in vivo behavior of the model complexes (2a-6a, 9a, 10a), we chose to evaluate the in vitro/in vivo biol. behavior of a new tumor-seeking Bombesin pyrazolyl conjugate, [(L6)-G-G-G-Q-W-A-V-G-H-L-M-NH<sub>2</sub>], that has been labeled with the [<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> metal fragment. Stability, in vitro cell binding assays, and pharmacokinetics studies in normal mice are reported herein.

CC 8-9 (Radiation Biochemistry)

Section cross-reference(s): 63, 78

- ST pyrazolyl bifunctional chelator technetium 99m prepn  
biodistribution; rhenium pyrazolyl complex prepn
- IT Drug delivery systems  
(carriers; pyrazolyl derivs. as bifunctional chelators for  
labeling tumor-seeking peptides with 99mTc or Re)
- IT Human  
Imaging agents  
Neoplasm  
Prostate gland, neoplasm  
Radiopharmaceuticals  
(pyrazolyl derivs. as bifunctional chelators for labeling  
tumor-seeking peptides with 99mTc or Re)
- IT 795271-47-5P 796031-75-9P 850494-15-4P 850494-16-5P 850494-17-6P  
850494-18-7P 850494-19-8P 850494-21-2P 850494-22-3P  
RL: DGN (Diagnostic use); PKT (Pharmacokinetics); PRP (Properties); SPN  
(Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES  
(Uses)  
(pyrazolyl derivs. as bifunctional chelators for labeling  
tumor-seeking peptides with 99mTc or Re)
- IT 511513-19-2P 511513-21-6P 511513-22-7P 782501-83-1P 782501-84-2P  
782501-85-3P 850494-14-3P 850494-20-1P  
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
(pyrazolyl derivs. as bifunctional chelators for labeling  
tumor-seeking peptides with 99mTc or Re)
- IT 107-15-3, Ethylenediamine, reactions 2087-41-4, Glycylglycine ethyl  
ester hydrochloride 2969-81-5, Ethyl 4-bromobutyrate 25908-22-9  
119291-22-4, 1-(2-Bromoethyl)pyrazole 163932-31-8 511513-23-8  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(pyrazolyl derivs. as bifunctional chelators for labeling  
tumor-seeking peptides with 99mTc or Re)
- IT 782501-70-6P 782501-71-7P 782501-72-8P  
782501-76-2P 782501-77-3P 782501-78-4P  
782501-79-5P 782501-80-8P 850480-64-7P  
850480-65-8P 850480-66-9P 850480-67-0P  
850480-68-1P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
(Reactant or reagent)  
(pyrazolyl derivs. as bifunctional chelators for labeling  
tumor-seeking peptides with 99mTc or Re)
- IT 511513-23-8  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(pyrazolyl derivs. as bifunctional chelators for labeling  
tumor-seeking peptides with 99mTc or Re)
- RN 511513-23-8 HCAPLUS
- CN 1,2-Ethanediamine, N1-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]- (CA INDEX  
NAME)



- IT 782501-70-6P 782501-71-7P 782501-72-8P  
782501-76-2P 782501-77-3P 782501-78-4P  
782501-79-5P 782501-80-8P 850480-64-7P

10/551,292

850480-65-8F 850480-66-9F 850480-67-0F


850480-68-1F

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(pyrazolyl derivs. as bifunctional chelators for labeling tumor-seeking peptides with 99mTc or Re)


RN 782501-70-6 HCAPLUS

CN 1,2-Ethanediamine, N1-[2-(1H-pyrazol-1-yl)ethyl]- (CA INDEX NAME)




RN 782501-71-7 HCAPLUS

CN Carbamic acid, [2-[[2-(1H-pyrazol-1-yl)ethyl]amino]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)




RN 782501-72-8 HCAPLUS

CN Butanoic acid, 4-[(2-aminoethyl)[2-(1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



RN 782501-76-2 HCAPLUS

CN Carbamic acid, N-[2-[[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]ethyl]-, 1,1-dimethylethyl ester (CA INDEX NAME)




RN 782501-77-3 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-


10/551,292

dimethyl-1H-pyrazol-1-yl)ethyl]amino]-, ethyl ester (CA INDEX NAME)




RN 782501-78-4 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-(CH2)3]butanoic acid (CA INDEX NAME)




RN 782501-79-5 HCAPLUS

CN 2,5,10,13-Tetraazapentadecanedioic acid, 5-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-9,12-dioxo-, 1-[(1,1-dimethylethyl) 15-ethyl ester (CA INDEX NAME)




RN 782501-80-8 HCAPLUS

CN Glycine, N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]glycyl-, ethyl ester (CA INDEX NAME)




RN 850480-64-7 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(1H-pyrazol-1-yl)ethyl]amino]-, ethyl ester (CA INDEX NAME)




RN 850480-65-8 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)




RN 850480-66-9 HCAPLUS

CN Butanoic acid, 4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)




RN 850480-67-0 HCAPLUS

CN 2,5,10,13-Tetraazapentadecanedioic acid, 9,12-dioxo-5-[2-(1H-pyrazol-1-yl)ethyl]-, 1-(1,1-dimethylethyl) 15-ethyl ester (CA INDEX NAME)



RN 850480-68-1 HCAPLUS

CN Glycine, N-[4-[(2-aminoethyl)[2-(1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]glycyl-, ethyl ester (CA INDEX NAME)



REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 9 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 9

ACCESSION NUMBER: 2004:902222 HCAPLUS Full-text

DOCUMENT NUMBER: 141:387794

TITLE: Preparation of bifunctional pyrazole-containing tridentate ligands for rhenium and technetium tricarbonyl complexes

INVENTOR(S): Santos, Isabel R.; Galamba Correia, Joao D.; Rocha Paulo, Antonio M.; Alves, Susana; Vitor, Rute

PATENT ASSIGNEE(S): Mallinckrodt Inc., USA

SOURCE: PCT Int. Appl., 60 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004091669	A1	20041028	WO 2004-US11685	20040415
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1529537	A1	20050511	EP 2003-78217	20031010
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
AU 2004229568	A1	20041028	AU 2004-229568	20040415
CA 2522326	A1	20041028	CA 2004-2522326	20040415
EP 1644050	A1	20060412	EP 2004-759566	20040415
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK			
CN 1774268	A	20060517	CN 2004-80010214	20040415
JP 2007525452	T	20070906	JP 2006-510091	20040415
US 20060198785	A1	20060907	US 2005-551292	20050928
IN 2005CN02650	A	20070525	IN 2005-CN2650	20051014
NO 2005005334	A	20051111	NO 2005-5334	20051111
PRIORITY APPLN. INFO.:			EP 2003-76106	A 20030415

OTHER SOURCE(S): MARPAT 141:387794  
ED Entered STN: 28 Oct 2004  
GI



AB The present invention relates to a chelating agent I [m = 0, 1; X = NR<sub>4</sub>, S; Y = SR<sub>5</sub>, NHR<sub>5</sub>, P(R<sub>5</sub>)<sub>2</sub>; R<sub>1</sub>, R<sub>3</sub> = independently H, alkyl, aryl; R<sub>2</sub> = H, CO<sub>2</sub>H, NHR<sub>6</sub>, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sub>6</sub>; R<sub>4</sub> = H, alkyl, aryl, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sub>6</sub>, (CH<sub>2</sub>)<sub>n</sub>OR<sub>6</sub>; R<sub>5</sub> = H, alkyl, aryl, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sub>6</sub>, (CH<sub>2</sub>)<sub>n</sub>OR<sub>6</sub>, R<sub>6</sub> = H, alkyl, aryl; n = 1-10; when R<sub>1</sub> = R<sub>3</sub> = CH<sub>3</sub>, R<sub>2</sub>, R<sub>4</sub>, R<sub>5</sub> are not all = H]. The invention further relates to a method and kit for the preparation of radiolabeled biomols. while using the chelating agent. Thus, pyrazole II (R = CO<sub>2</sub>H) was prepared by cyclocondensation of (OHC)<sub>2</sub>CHCO<sub>2</sub>Et with H<sub>2</sub>NNHCH<sub>2</sub>CH<sub>2</sub>OH, followed by tosylation and substitution with ethylenediamine and saponification Prepared compds. II (R = H, CO<sub>2</sub>H) underwent complexation with rhenium and technetium-99 to give the corresponding tricarbonyl complexes.

IC ICM A61K051-04

ICS C07D231-12; C07D231-14; A61P035-00

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 9, 28

IT Chelating agents

(preparation of bifunctional pyrazole-containing tridentate ligands for rhenium

and technetium tricarbonyl complexes)

IT 60-24-2, 2-Mercaptoethanol 107-15-3, Ethylenediamine, reactions  
109-84-2, (2-Hydroxyethyl)hydrazine 623-51-8, Ethyl 2-mercaptoacetate  
2087-41-4, Glycylglycine ethyl ester hydrochloride 2969-81-5, Ethyl  
4-bromobutyrate 67000-34-4 80370-42-9 119291-22-4,  
1-(2-Bromoethyl)pyrazole 511513-23-8

RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of bifunctional pyrazole-containing tridentate ligands for rhenium

and technetium tricarbonyl complexes)

IT 487021-85-2P 782501-70-6P 782501-71-7P 782501-73-9P  
782501-74-0P 782501-75-1P 782501-76-2P  
782501-77-3P 782501-78-4P 782501-79-5P  
782501-80-8P 782501-81-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of bifunctional pyrazole-containing tridentate ligands for rhenium

and technetium tricarbonyl complexes)

IT 782501-72-8P 782501-82-0P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of bifunctional pyrazole-containing tridentate ligands for rhenium


and technetium tricarbonyl complexes)

IT 511513-23-8




10/551,292

RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation of bifunctional pyrazole-containing tridentate ligands for  
 rhenium and technetium tricarbonyl complexes)  
 RN 511513-23-8 HCAPLUS  
 CN 1,2-Ethanediamine, N1-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]- (CA INDEX  
 NAME)




IT 782501-70-6P 782501-71-7P 782501-75-1P  
782501-76-2P 782501-77-3P 782501-78-4P  
782501-79-5P 782501-80-8P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preparation of bifunctional pyrazole-containing tridentate ligands for  
 rhenium and technetium tricarbonyl complexes)  
 RN 782501-70-6 HCAPLUS  
 CN 1,2-Ethanediamine, N1-[2-(1H-pyrazol-1-yl)ethyl]- (CA INDEX NAME)



RN 782501-71-7 HCAPLUS  
 CN Carbamic acid, [2-[[2-(1H-pyrazol-1-yl)ethyl]amino]ethyl]-,  
 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)




RN 782501-75-1 HCAPLUS  
 CN 1H-Pyrazole-4-carboxylic acid, 1-[2-[(2-aminoethyl)amino]ethyl]- (CA  
 INDEX NAME)




RN 782501-76-2 HCAPLUS

CN Carbamic acid, N-[2-[[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]ethyl]-, 1,1-dimethylethyl ester (CA INDEX NAME)




RN 782501-77-3 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-, ethyl ester (CA INDEX NAME)




RN 782501-78-4 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)




RN 782501-79-5 HCAPLUS

CN 2,5,10,13-Tetraazapentadecanedioic acid, 5-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-9,12-dioxo-, 1-(1,1-dimethylethyl) 15-ethyl ester (CA INDEX NAME)



RN 782501-80-8 HCAPLUS

CN Glycine, N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]glycyl-, ethyl ester (CA INDEX NAME)




IT 782501-72-8P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of bifunctional pyrazole-containing tridentate ligands for rhenium and technetium tricarbonyl complexes)

RN 782501-72-8 HCAPLUS

CN Butanoic acid, 4-[(2-aminoethyl)[2-(1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 10 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 10

ACCESSION NUMBER: 2004:1049014 HCAPLUS Full-text

DOCUMENT NUMBER: 142:168330

TITLE: Rhenium(I)- and technetium(I) tricarbonyl complexes anchored by bifunctional pyrazole-diamine and pyrazole-dithioether chelators

AUTHOR(S): Vitor, Rute F.; Alves, Susana; Correia, J. D. G.; Paulo, Antonio; Santos, Isabel

CORPORATE SOURCE: ITN, Estrada Nacional, Departamento de Quimica, Sacavem Codex, 2686-953, Port.

SOURCE: Journal of Organometallic Chemistry (2004), 689(25), 4764-4774

CODEN: JORCAI; ISSN: 0022-328X

PUBLISHER: Elsevier B.V.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 142:168330

ED Entered STN: 08 Dec 2004

AB The novel pyrazolyl containing ligands 4-(HOOC)pz(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub> (L1) and 4-(HOOCCH<sub>2</sub>)-3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub> (L2), and 3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>S(CH<sub>2</sub>)<sub>2</sub>SCH<sub>2</sub>CH<sub>3</sub> (L3), 3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>S(CH<sub>2</sub>)<sub>2</sub>SCH<sub>2</sub>COOEt (L4) and 3,5-Me<sub>2</sub>pz(CH<sub>2</sub>)<sub>2</sub>S(CH<sub>2</sub>)<sub>2</sub>SCH<sub>2</sub>COOH (L5) were synthesized, and their ability to stabilize complexes with the fac-[M(CO)<sub>3</sub>]<sup>+</sup> (M = Re, <sup>99m</sup>Tc) moiety was evaluated. Reactions of L1-L5 with (NEt<sub>4</sub>)<sub>2</sub>[Re(CO)<sub>3</sub>Br<sub>3</sub>] and/or [Re(CO)<sub>5</sub>Br] afforded complexes fac-[Re(CO)<sub>3</sub>(κ<sup>3</sup>-L)] (L = L1-L5 (1-5)), which contain the pyrazolyl ancillary ligands coordinated in a tridentate fashion. Complexes 1-5 were characterized by the common anal. techniques, which included single crystal x-ray diffraction anal. in the case of 4. The structural anal. of 4 confirmed the tridentate coordination mode of the pyrazole-dithioether ligand, which is facially coordinated to the Re(I) center through the N from the pyrazole ring and the two thioether S atoms, without involvement of the terminal ester functional group. The distorted octahedral coordination environment around the metal is completed by the three facial carbonyl ligands. The radioactive congeners of complexes 1, 3 and 4, fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>(κ<sup>3</sup>-L)]<sup>+</sup> (L = L1 (1a), L3 (3a), L4 (4a)), were prepared by reacting the precursor fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>(H<sub>2</sub>O)<sub>3</sub>]<sup>+</sup> with the corresponding ligands, and their identity confirmed by HPLC comparison with the Re surrogates. Complexes 1a and 3a were challenged in the presence of a large excess of histidine or cysteine, to evaluate their in vitro stability. Only a negligible displacement was observed, indicating that pyrazole-diamine and pyrazole-dithioether chelators provide a high kinetic inertness and/or stability to organometallic complexes with the fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> moiety.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 28, 75

ST rhenium carbonyl pyrazole dithioether diamine chelator complex  
 prepn; crystal structure rhenium carbonyl pyrazole dithioether  
chelator complex; stability in vitro technetium pyrazole  
 dithioether chelator complex; technetium pyrazole dithioether  
 diamine chelator complex prepn

IT Stability

(stability of technetium carbonyl pyrazole-thioether and -diamine  
chelator complexes in presence of histidine and cysteine)

IT 163932-31-8, fac-Triaquatricarbonyltechnetium(1+)-<sup>99</sup>Tc

RL: RCT (Reactant); RACT (Reactant or reagent)

(metastable; reactant for preparation of technetium carbonyl  
 pyrazole-thioether and -diamine chelator complexes)

IT 827596-91-8P 827596-93-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)

(preparation and complexation with rhenium)

IT 782501-82-0P 827596-90-7P 827596-92-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)

(preparation and complexation with rhenium and technetium)

IT 782501-73-9P 827596-95-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)

(preparation and reactant for preparation of pyrazole-diamine chelator)

IT 827596-94-1P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)

(preparation and reactant for preparation of pyrazole-diamine chelators  
 )

IT 782501-81-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and reactant for preparation of pyrazole-thioether chelators)

IT 1656-44-6, 2,4-Dinitrobenzenesulfonyl chloride 57260-73-8,  
N-tert-Butoxycarbonyl-1,2-ethanediamine 80370-42-9 503471-30-5

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant for preparation of pyrazole-diamine chelator)

IT 109-84-2, 2-Hydroxyethylhydrazine

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant for preparation of pyrazole-diamine chelators)

IT 75-08-1, Ethanethiol 623-51-8, Ethyl 2-mercaptoacetate 487021-85-2

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant for preparation of pyrazole-thioether chelators)

IT 14220-21-4, Bromopentacarbonylrhenium

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant for preparation of rhenium carbonyl pyrazole-diamine chelator complexes)

IT 25908-22-9, Bis(tetraethylammonium) tribromotricarbonylrhenate(2-)

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant for preparation of rhenium carbonyl pyrazole-thioether chelator complexes)

IT 52-90-4, L-Cysteine, processes 71-00-1, L-Histidine, processes

RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PROC (Process)

(stability of technetium carbonyl pyrazole-thioether and -diamine chelator complexes in presence of)


IT 827596-91-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and complexation with rhenium)

RN 827596-91-8 HCAPLUS

CN 1H-Pyrazole-4-acetic acid, 1-[2-[(2-aminoethyl)amino]ethyl]-3,5-dimethyl-, ethyl ester (CA INDEX NAME)




IT 827596-90-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and complexation with rhenium and technetium)

RN 827596-90-7 HCAPLUS

CN 1H-Pyrazole-4-carboxylic acid, 1-[2-[(2-aminoethyl)amino]ethyl]-, ethyl ester (CA INDEX NAME)



REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 11 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:1487958 HCAPLUS Full-text

DOCUMENT NUMBER: 150:208111

TITLE: Re and 99m Tc organometallic complexes containing pendant L-arginine derivatives as potential probes of inducible nitric oxide synthase

AUTHOR(S): Oliveira, Bruno L.; Correia, Joao D. G.;

Raposo, Paula D.; Santos, Isabel;

Ferreira, Antonio; Cordeiro, Carlos; Freire, Ana P.

CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem, 2686-953, Port.

SOURCE: Dalton Transactions (2009), (1), 152-162

CODEN: DTARAF; ISSN: 1477-9226

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 12 Dec 2008

AB Aiming to design radioactive compds. based on the core "99mTc(CO)3" for probing inducible nitric oxide synthase (iNOS) levels in vivo, we have synthesized conjugates containing a pyrazolyl-diamine chelating unit and pendant L-arginine analogs (substrates and inhibitors of NOS). Reaction of the conjugates with fac-[M(CO)3]+ (M = Re, 99m Tc) gave bioorganometallic complexes of the type fac-[M(CO)3(k3-L)] in good yield. After in vitro testing using the oxyHb NO capture assay, we concluded that the affinity of the inhibitor-containing conjugates to iNOS seems to be less affected upon metalation with rhenium than the substrate-containing conjugates. The complexes bearing guanidino substituted analogs of L-arginine still present considerable inhibitory action (N $\omega$ -monomethyl-L-arginine, Ki = 36  $\mu$ M; N $\omega$ -nitro-L-arginine, Ki = 84  $\mu$ M), being the first examples of organometallic complexes able to inhibit the iNOS. These results seem to indicate that 99mTc(CO)3-labeled L-arginine analogs, namely NOS inhibitors, may hold potential for monitoring increased levels of iNOS in vivo.

CC 7-3 (Enzymes)

Section cross-reference(s): 8, 9, 78

IT 74-79-3, L-Arginine, reactions 2577-94-8, L-Arginine methyl ester

6066-82-6, N-Hydroxysuccinimide 17035-90-4 25908-22-9,

Bis(tetraethylammonium) fac-tribromotricarbonylrhenate(2-) 50903-99-6,

N $\omega$ -Nitro-L-arginine methyl ester 163932-31-8,

Fac-triaquatricarbonyltechnetium-99(1+) 850480-66-9

945384-90-7 1111224-57-7 1111224-64-6

1111224-73-7

RL: RCT (Reactant); RACT (Reactant or reagent)

(Re and 99mTc organometallic complexes containing pendant L-arginine derivs. as potential probes of inducible nitric oxide synthase)

IT 1111224-18-0P 1111224-21-5P 1111224-23-7P

1111224-25-9P 1111224-29-3P 1111224-32-8P

1111224-35-1P 1111224-37-3P 1111224-40-8P

1111224-43-1P 1111224-46-4P 1111224-48-6P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(Re and 99mTc organometallic complexes containing pendant L-arginine derivs. as potential probes of inducible nitric oxide synthase)

IT 850480-66-9 1111224-57-7 1111224-64-6

1111224-73-7


10/551,292

RL: RCT (Reactant); RACT (Reactant or reagent)

(Re and 99mTc organometallic complexes containing pendant L-arginine derivs. as potential probes of inducible nitric oxide synthase)

RN 850480-66-9 HCAPLUS

CN Butanoic acid, 4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



RN 1111224-57-7 HCAPLUS


CN L-Ornithine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-N5-[imino(nitroamino)methyl]-, methyl ester, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1

CRN 1111224-40-8

CMF C20 H37 N9 O5


Absolute stereochemistry.



CM 2

CRN 76-05-1

CMF C2 H F3 O2



RN 1111224-64-6 HCAPLUS

CN L-Ornithine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-N5-[imino(nitroamino)methyl]-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)


10/551,292

CM 1

CRN 1111224-43-1

CMF C19 H35 N9 O5


Absolute stereochemistry.



CM 2

CRN 76-05-1

CMF C2 H F3 O2



RN 1111224-73-7 HCAPLUS


CN L-Ornithine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-N5-[imino(methylamino)methyl]-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1

CRN 1111224-48-6

CMF C20 H38 N8 O3

Absolute stereochemistry.




CM 2

CRN 76-05-1

CMF C2 H F3 O2






IT 1111224-18-0P 1111224-21-5P 1111224-23-7P  
1111224-25-9P 1111224-29-3P 1111224-32-8P  
1111224-35-1P 1111224-37-3P 1111224-40-8P  
1111224-43-1P 1111224-46-4P 1111224-48-6P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(Re and <sup>99m</sup>Tc organometallic complexes containing pendant L-arginine derivs. as potential probes of inducible nitric oxide synthase)

RN 1111224-18-0 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-, 2,5-dioxo-1-pyrrolidinyl ester (CA INDEX NAME)



RN 1111224-21-5 HCAPLUS


CN 13-Oxa-3,8,11-triazapentadecanoic acid, 2-[3-[(aminoiminomethyl)amino]propyl]-8-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-14,14-dimethyl-4,12-dioxo-, methyl ester, (2S)-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1

CRN 1111224-20-4

CMF C25 H46 N8 O5

Absolute stereochemistry.




10/551,292

CM 2

CRN 76-05-1


CMF C2 H F3 O2



RN 1111224-23-7 HCAPLUS

CN L-Arginine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-, methyl ester (CA INDEX NAME)

Absolute stereochemistry.



RN 1111224-25-9 HCAPLUS


CN L-Arginine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-, methyl ester, 2,2,2-trifluoroacetate (1:2) (CA INDEX NAME)

CM 1

CRN 1111224-23-7

CMF C20 H38 N8 O3

Absolute stereochemistry.




CM 2

CRN 76-05-1

CMF C2 H F3 O2

10/551,292




RN 1111224-29-3 HCAPLUS  
 CN 13-Oxa-3,8,11-triazapentadecanoic acid,  
 2-[3-[(aminoiminomethyl)amino]propyl]-8-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-14,14-dimethyl-4,12-dioxo-, (2S)-, 2,2,2-trifluoroacetate (1:1)  
 (CA INDEX NAME)

CM 1


CRN 1111224-28-2  
 CMF C24 H44 N8 O5

Absolute stereochemistry.



CM 2


CRN 76-05-1  
 CMF C2 H F3 O2



RN 1111224-32-8 HCAPLUS  
 CN L-Arginine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]- (CA INDEX NAME)

Absolute stereochemistry.

10/551,292



RN 1111224-35-1 HCAPLUS


CN L-Arginine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1

CRN 1111224-32-8

CMF C19 H36 N8 O3


Absolute stereochemistry.



CM 2

CRN 76-05-1

CMF C2 H F3 O2




RN 1111224-37-3 HCAPLUS

CN 2,5,10,15-Tetraazahexadecanoic acid, 5-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-16-imino-11-(methoxycarbonyl)-16-(nitroamino)-9-oxo-, 1,1-dimethylethyl ester, (11S)- (CA INDEX NAME)

Absolute stereochemistry.


10/551,292



RN 1111224-40-8 HCAPLUS

CN L-Ornithine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-N5-[imino(nitroamino)methyl]-, methyl ester  
(CA INDEX NAME)


Absolute stereochemistry.



RN 1111224-43-1 HCAPLUS

CN L-Ornithine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-N5-[imino(nitroamino)methyl]- (CA INDEX NAME)


Absolute stereochemistry.



RN 1111224-46-4 HCAPLUS


CN 13-Oxa-3,8,11-triazapentadecanoic acid,  
8-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-2-[3-  
[[imino(methylamino)methyl]amino]propyl]-14,14-dimethyl-4,12-dioxo-, (2S)-  
(CA INDEX NAME)

Absolute stereochemistry.



RN 1111224-48-6 HCAPLUS  
 CN L-Ornithine, N2-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-N5-[imino(methylamino)methyl]- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 12 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2008:1206988 HCAPLUS [Full-text](#)  
 DOCUMENT NUMBER: 150:35664  
 TITLE: Synthesis, characterization, and evaluation of a novel 99mTc(CO)3 pyrazolyl conjugate of a peptide nucleic acid sequence. [Erratum to document cited in CA150:020350]  
 AUTHOR(S): Xavier, Catarina; Giannini, Clelia; Dall'Angelo, Sergio; Gano, Lurdes; Maiorana, Stefano; Alberto, Roger; Santos, Isabel  
 CORPORATE SOURCE: Departamento de Quimica, Instituto Tecnologico e Nuclear, Sacavem, 2686-953, Port.  
 SOURCE: JBIC, Journal of Biological Inorganic Chemistry (2008), 13(8), 1345  
 CODEN: JJBCFA; ISSN: 0949-8257  
 PUBLISHER: Springer GmbH  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 ED Entered STN: 08 Oct 2008  
 AB On page 1335, in the author list, Sergio Dall'Angelo was omitted from the author list; the correct author list and author affiliations are given.  
 CC 34-3 (Amino Acids, Peptides, and Proteins)  
 Section cross-reference(s): 8, 78  
 IT Nucleic acid hybridization  
 (DNA-PNA; solid-phase preparation of peptide nucleic acid sequences, their conjugates with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))  
 IT Imaging agents

(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))

IT Peptide nucleic acids  
RL: BSU (Biological study, unclassified); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)  
(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))

IT 1089234-29-6P  
RL: BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))

IT 1089234-31-0 1089234-32-1  
RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)  
(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))

IT 1089234-26-3P 1089234-28-5P  
RL: PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)  
(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))

IT 1089234-27-4P  
RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))


IT 163932-31-8 782501-78-4 945384-90-7  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))

IT 1089234-30-9DP, resin-bound  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))

IT 782501-78-4  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(solid-phase preparation of peptide nucleic acid sequences, their conjugates  
with pyrazolyl chelator, and complexes with rhenium and 99-technetium (Erratum))

RN 782501-78-4 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



L49 ANSWER 13 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:462227 HCAPLUS Full-text

DOCUMENT NUMBER: 148:556216

TITLE: Melanoma targeting with  $\alpha$ -melanocyte stimulating hormone analogs labeled with fac-[ $^{99m}\text{Tc}(\text{CO})_3$ ] $^{+}$ : effect of cyclization on tumor-seeking properties

AUTHOR(S): Raposinho, Paula D.; Xavier, Catarina; Correia, Joao D. G.; Falcao, Soraia; Gomes, Paula; Santos, Isabel

CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem, 2686-953, Port.

SOURCE: JBIC, Journal of Biological Inorganic Chemistry

(2008), 13(3), 449-459

CODEN: JJBCFA; ISSN: 0949-8257

PUBLISHER: Springer GmbH

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 15 Apr 2008

AB Early detection of primary melanoma tumors is essential because there is no effective treatment for metastatic melanoma. Several linear and cyclic radiolabeled  $\alpha$ -MSH analogs have been proposed to target the melanocortin type 1 receptor (MC1R) overexpressed in melanoma. The compact structure of a rhenium-cyclized  $\alpha$ -MSH analog (Re-CCMSH) significantly enhanced its in vivo tumor uptake and retention. Melanotan II (MT-II), a cyclic lactam analog of  $\alpha$ -MSH (Ac-Nle-cyclo[Asp-His-DPhe-Arg-Trp-Lys]-NH<sub>2</sub>), is a very potent and stable agonist peptide largely used in the characterization of melanocortin receptors. Taking advantage of the superior biol. features associated with the MT-II cyclic peptide, the authors assessed the effect of lactam-based cyclization on the tumor-seeking properties of  $\alpha$ -MSH analogs by comparing the pharmacokinetics profile of the  $^{99m}\text{Tc}$ -labeled cyclic peptide  $\beta\text{Ala-Nle-cyclo[Asp-His-D-Phe-Arg-Trp-Lys]-NH}_2$  with that of the linear analog  $\beta\text{Ala-Nle-Asp-His-DPhe-Arg-Trp-Lys-NH}_2$  in melanoma-bearing mice. The authors have synthesized and coupled the linear and cyclic peptides to a bifunctional chelator containing a pyrazolyl-diamine backbone (pz) through the amino group of  $\beta\text{Ala}$ , and the resulting pz-peptide conjugates were reacted with the fac-[ $^{99m}\text{Tc}(\text{CO})_3$ ] $^{+}$  moiety. The  $^{99m}\text{Tc}(\text{CO})_3$ -labeled conjugates were obtained in high yield, high specific activity, and high radiochem. purity. The cyclic  $^{99m}\text{Tc}(\text{CO})_3$ -labeled conjugate presents a remarkable internalization (87.1% of receptor-bound tracer and 50.5% of total applied activity, after 6 h at 37°) and cellular retention (only 24.7% released from the cells after 5 h) in murine melanoma B16F1 cells. A significant tumor uptake and retention was obtained in melanoma-bearing C57BL6 mice for the cyclic radioconjugate [ $9.26 \pm 0.83$  and  $11.31 \pm 1.83\%$  ID/g at 1 and 4 h after injection, resp.]. The linear  $^{99m}\text{Tc}(\text{CO})_3$ -pz-peptide presented lower values for both cellular internalization



and tumor uptake. Receptor blocking studies with the potent (Nle<sup>4</sup>,DPhe<sup>7</sup>)- $\alpha$ MSH agonist demonstrated the specificity of the radio-conjugates to MC1R (74.8% and 44.5% reduction of tumor uptake at 4 h after injection for cyclic and linear radio-conjugates, resp.).

CC 8-9 (Radiation Biochemistry)

Section cross-reference(s): 2, 14

IT 524744-56-7 782501-78-4

RL: RCT (Reactant); RACT (Reactant or reagent)

(melanoma targeting with  $\alpha$ -MSH analogs labeled with fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> and effect of cyclization on tumor-seeking properties)

IT 1025483-81-1P 1025483-83-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(melanoma targeting with  $\alpha$ -MSH analogs labeled with fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> and effect of cyclization on tumor-seeking properties)


IT 782501-78-4

RL: RCT (Reactant); RACT (Reactant or reagent)

(melanoma targeting with  $\alpha$ -MSH analogs labeled with fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> and effect of cyclization on tumor-seeking properties)

RN 782501-78-4 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



IT 1025483-81-1P 1025483-83-3P




RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(melanoma targeting with  $\alpha$ -MSH analogs labeled with fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> and effect of cyclization on tumor-seeking properties)

RN 1025483-81-1 HCAPLUS

CN L-Lysinamide, N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]- $\beta$ -alanyl-L-norleucyl-L- $\alpha$ -aspartyl-L-histidyl-D-phenylalanyl-L-arginyl-L-tryptophyl-, (3 $\rightarrow$ 8)-lactam (CA INDEX NAME)



Absolute stereochemistry.



RN 1025483-83-3 HCAPLUS


CN L-Lysinamide, N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-β-alanyl-L-norleucyl-L-α-aspartyl-L-histidyl-D-phenylalanyl-L-arginyl-L-tryptophyl- (CA INDEX NAME)

Absolute stereochemistry.






REFERENCE COUNT: 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 14 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2008:1176860 HCAPLUS Full-text  
 DOCUMENT NUMBER: 149:576827  
 TITLE: Comparative Study of Chemical Approaches to the Solid-Phase Synthesis of a Tumor-Seeking  $\alpha$ -MSH Analogue  
 AUTHOR(S): Valldosera, Magdalena; Monso, Marta; Xavier, Catarina; Raposinho, Paula; Correia, Joao D. G.; Santos, Isabel; Gomes, Paula  
 CORPORATE SOURCE: Departamento de Quimica, Faculdade de Ciencias (DQFCUP), Centro de Investigacao em Quimica da Universidade do Porto (CIQUP), Universidade do Porto, Oporto, 4169-007, Port.  
 SOURCE: International Journal of Peptide Research and Therapeutics (2008), 14(3), 273-281  
 CODEN: IJPRFC; ISSN: 1573-3149  
 PUBLISHER: Springer  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 ED Entered STN: 01 Oct 2008  
 GI



- AB The synthesis of a cyclic melanocortin analog I (H-pz- $\beta$ Ala-Nle-cyclo[Asp-His-DPhe-Arg-Trp-Lys]-NH<sub>2</sub>), where the Boc-protected derivative of a metal-chelating pyrazolyl ligand (pz) was inserted as an N-terminal residue, was accomplished by several different Fmoc/tBu and Boc/Bzl solid-phase strategies. On-resin cyclization was achieved immediately following incorporation of Asp, by condensation of the Asp side chain carboxyl with the Lys side chain primary amine after selective and simultaneous removal of side chain protecting groups. The success of the synthesis was highly dependent on the chemical strategy employed, with Boc/Bzl chemical giving the best results. On the light of these findings, Fmoc/tBu strategies are not advantageous for the solid-phase synthesis of this particular type of lactam-bridged peptides. Last, but not least, the target peptide was recently found to have promising tumor-seeking properties (J Biol Inorg Chem 13:449-459, 2008).
- CC 34-3 (Amino Acids, Peptides, and Proteins)  
Section cross-reference(s): 2
- IT 1025483-81-1P  
RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(comparisons of solid-phase synthetic approaches to tumor-seeking  $\alpha$ -MSH analog)
- IT 81379-52-4 84624-27-1 109425-51-6 117014-32-1 146982-24-3  
146982-27-6 167393-62-6 200336-86-3 204777-78-6, Fmoc-Lys(ivDde)-OH  
269066-08-2 782501-78-4  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(comparisons of solid-phase synthetic approaches to tumor-seeking  $\alpha$ -MSH analog)
- IT 137668-62-3P 1084652-81-2P 1084652-83-4P 1084652-84-5P  
1084652-87-8P 1084652-90-3P 1084652-93-6P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(comparisons of solid-phase synthetic approaches to tumor-seeking  $\alpha$ -MSH analog)
- IT 1025483-81-1P  
RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(comparisons of solid-phase synthetic approaches to tumor-seeking  $\alpha$ -MSH analog)
- RN 1025483-81-1 HCAPLUS
- CN L-Lysinamide, N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]- $\beta$ -alanyl-L-norleucyl-L- $\alpha$ -aspartyl-L-histidyl-D-phenylalanyl-L-arginyl-L-tryptophyl-, (3 $\rightarrow$ 8)-lactam (CA INDEX NAME)

Absolute stereochemistry.


IT 782501-78-4

RL: RCT (Reactant); RACT (Reactant or reagent)

(comparisons of solid-phase synthetic approaches to tumor-seeking  
 $\alpha$ -MSH analog)

RN 782501-78-4 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



IT 1084652-81-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(comparisons of solid-phase synthetic approaches to tumor-seeking  $\alpha$ -MSH analog)



RN 1084652-81-2 HCAPLUS

CN L-Lysinamide, N-[4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]- $\beta$ -alanyl-L-norleucyl-L- $\alpha$ -aspartyl-L-histidyl-D-phenylalanyl-L-arginyl-L-tryptophyl-N6-[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl]-, [4-[[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl]amino]phenyl]methyl ester (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 15 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:605111 HCAPLUS Full-text

DOCUMENT NUMBER: 147:229603

TITLE: A new bisphosphonate-containing <sup>99m</sup>Tc(I) tricarbonyl complex potentially useful as bone-seeking agent: synthesis and biological evaluation

AUTHOR(S): Palma, Elisa; Oliveira, Bruno L.; Correia, Joao D. G.; Gano, Lurdes; Maria, Leonor; Santos, Isabel C.; Santos, Isabel

CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem Codex, 2686-953, Port.

SOURCE: JBIC, Journal of Biological Inorganic Chemistry (2007), 12(5), 667-679  
CODEN: JJBCFA; ISSN: 0949-8257

PUBLISHER: Springer GmbH

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 05 Jun 2007

AB Aiming to develop new bone-seeking radiotracers based on the organometallic core fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>]<sup>+</sup> with improved radiochem. and biol. properties, we have prepared new conjugates with phosphonate pendant groups. The conjugates comprise a chelating unit for metal coordination, which corresponds to a pyrazolyl-containing backbone (pz) with a N,N,N donor-atom set, and a pendant

di-Et phosphonate (pz-MPOEt), phosphonic acid (pz-MPOH) or a bisphosphonic acid (pz-BPOH) group for bone targeting. Reactions of the conjugates with the precursor  $[^{99m}\text{Tc}(\text{H}_2\text{O})_3(\text{CO})_3]^+$  yielded (more than 95%) the single and well-defined radioactive species  $[^{99m}\text{Tc}(\text{CO})_3(\kappa^3\text{-pz-MPOEt})]^+$  (1a),  $[^{99m}\text{Tc}(\text{CO})_3(\kappa^3\text{-pz-MPOH})]^+$  (2a) and  $[^{99m}\text{Tc}(\text{CO})_3(\kappa^3\text{-pz-BPOH})]^+$  (3a), which were characterized by reversed-phase high-performance liquid chromatog. . The corresponding Re surrogates (1-3), characterized by the usual anal. techniques, including X-ray diffraction anal. in the case of 1, allowed for macroscopic identification of the radioactive conjugates. These radioactive complexes revealed high stability both in vitro (phosphate-buffered saline solution and human plasma) and in vivo, without any measurable decomposition. Biodistribution studies of the complexes in mice indicated a fast rate of blood clearance and high rate of total radioactivity excretion, occurring primarily through the renal-urinary pathway in the case of complex 3a. Despite presenting moderate bone uptake ( $3.04 \pm 0.47\%$  injected dose per g of organ, 4 h after injection), the high stability presented by 3a and its adequate in vivo pharmacokinetics encourages the search for new ligands with the same chelating unit and different bisphosphonic acid pendant arms.

CC 8-9 (Radiation Biochemistry)

IT 5324-30-1 14220-21-4 25908-22-9 80474-99-3 782501-76-2

782501-78-4 850480-66-9 945384-86-1

RL: RCT (Reactant); RACT (Reactant or reagent)

(bisphosphonate-containing  $^{99m}\text{Tc}(\text{I})$  tricarbonyl complex: preparation and potential as bone-seeking agent)

IT 945264-41-5P 945264-42-6P 945264-43-7P

945264-45-9P 945264-46-0P 945384-90-7P 945384-92-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(bisphosphonate-containing  $^{99m}\text{Tc}(\text{I})$  tricarbonyl complex: preparation and potential as bone-seeking agent)


IT 782501-76-2 782501-78-4 850480-66-9

RL: RCT (Reactant); RACT (Reactant or reagent)

(bisphosphonate-containing  $^{99m}\text{Tc}(\text{I})$  tricarbonyl complex: preparation and potential as bone-seeking agent)

RN 782501-76-2 HCAPLUS


CN Carbamic acid, N-[2-[[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]ethyl]-, 1,1-dimethylethyl ester (CA INDEX NAME)



RN 782501-78-4 HCAPLUS


CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)





RN 850480-66-9 HCAPLUS

CN Butanoic acid, 4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



IT 945264-41-5P 945264-42-6P 945264-43-7P


945264-45-9P 945264-46-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(bisphosphonate-containing  $^{99m}\text{Tc}$ (I) tricarbonyl complex: preparation and potential as bone-seeking agent)


RN 945264-41-5 HCAPLUS

CN 9-Oxa-2,5-diaza-8-phosphaundecanoic acid, 5-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-8-ethoxy-, 1,1-dimethylethyl ester, 8-oxide (CA INDEX NAME)




RN 945264-42-6 HCAPLUS

CN Phosphonic acid, P-[2-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]ethyl]-, diethyl ester (CA INDEX NAME)




RN 945264-43-7 HCAPLUS

CN Phosphonic acid, P-[2-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]ethyl]- (CA INDEX NAME)




RN 945264-45-9 HCAPLUS

CN 13-Oxa-2,5,10-triaza-12-phosphapentadecanoic acid, 11-(diethoxyphosphinyl)-5-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-12-ethoxy-9-oxo-, 1,1-dimethylethyl ester, 12-oxide (CA INDEX NAME)



RN 945264-46-0 HCAPLUS

CN Phosphonic acid, P,P'-[[[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]amino]methylene]bis- (CA INDEX NAME)



REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 16 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:48914 HCAPLUS [Full-text](#)

DOCUMENT NUMBER: 146:311529

TITLE: In Vitro and In Vivo Evaluation of a Novel  
99mTc(CO)3-Pyrazolyl Conjugate of  
cyclo-(Arg-Gly-Asp-D-Tyr-Lys)

AUTHOR(S): [Alves, Susana](#); [Correia, Joao D. G.](#)  
; Gano, Lurdes; Rold, Tammy L.; Prasanphanich, Adam;  
Haubner, Roland; Rupprich, Marco; Alberto, Roger;

Decristoforo, Clemens; Santos, Isabel;  
 Smith, Charles J.

CORPORATE SOURCE: Department of Radiology Department of Internal  
 Medicine and The Radiopharmaceutical Sciences  
 Institute, University of Missouri-Columbia School of  
 Medicine, Columbia, MO, 65211, USA

SOURCE: Bioconjugate Chemistry (2007), 18(2), 530-537  
 CODEN: BCCHEs; ISSN: 1043-1802

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 146:311529

ED Entered STN: 16 Jan 2007

AB Radiolabeled peptides containing the Arg-Gly-Asp amino acid sequence (single  
 letter code = RGD) have been studied extensively to target integrin receptors  
 upregulated on tumor cells and neovasculature. Integrins are cell surface  
 transmembrane glycoproteins that exist as  $\alpha\beta$  heterodimers. The  $\alpha v\beta 3$  integrin  
 is known to be overexpressed in many tumor types and is expressed at lower  
 levels in normal tissues. Furthermore,  $\alpha v\beta 3$  and  $\alpha v\beta 5$  subtypes are expressed in  
 neovasculature during angiogenesis. Thus, there is some impetus to image  
 angiogenesis and tumor formation in vivo using RGD-based peptide targeting  
 vectors. In this study, we report the design and development of a new cyclic  
 RGD analog cyclo-[Arg-Gly-Asp-D-Tyr-Lys(PZ)] (PZ = 3,5-Me2-  
 pz(CH2)2N((CH2)3COOH)(CH2)2NH2) that can be radiolabeled with the  
 [99mTc(CO)3(H2O)3]+ metal aquaion. Radiochem. evaluation of this new  
 conjugate in vitro indicated a facile radiosynthesis of the new 99mTc-RGD  
 conjugate with high radiolabeling yields ( $\geq 95\%$ ) and high specific activities.  
 In vitro internalization and blocking assays in  $\alpha v\beta 3$  receptor-pos., human M21  
 melanoma cancer cells showed the ability of this conjugate to target the  
 integrin receptor with high specificity and selectivity. In vivo  
 pharmacokinetic studies in normal CF-1 mice showed rapid clearance from blood  
 with excretion primarily via/through the renal-urinary system. In vivo  
 accumulation of radioactivity in mice bearing either  $\alpha v\beta 3$  receptor-pos. or  
 neg. human melanoma tumors showed receptor specific uptake of tracer with  
 accumulations of  $2.50 \pm 0.29$  and  $0.71 \pm 0.08\%$  ID/g in  $\alpha v\beta 3$  integrin pos. (M21)  
 and neg. (M21L) tumors at 1 h postinjection (p.i.), resp.

CC 8-9 (Radiation Biochemistry)


IT 163932-31-8 217099-14-4 782501-78-4  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (99mTc(CO)3-pyrazolyl conjugate of cyclo-(Arg-Gly-Asp-D-Tyr-Lys)  
 preparation  
 and targeting integrin receptors in melanoma)

IT 928406-90-0P 928406-91-1P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (99mTc(CO)3-pyrazolyl conjugate of cyclo-(Arg-Gly-Asp-D-Tyr-Lys)  
 preparation  
 and targeting integrin receptors in melanoma)

IT 782501-78-4  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (99mTc(CO)3-pyrazolyl conjugate of cyclo-(Arg-Gly-Asp-D-Tyr-Lys)  
 preparation  
 and targeting integrin receptors in melanoma)

RN 782501-78-4 HCAPLUS

CN Butanoic acid, 4-[[2-[[[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-  
 dimethyl-1H-pyrazol-1-yl)ethyl]amino]- (CA INDEX NAME)



IT ~~928406-90-0P~~ ~~928406-91-1P~~

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(99mTc(CO)<sub>3</sub>-pyrazolyl conjugate of cyclo-(Arg-Gly-Asp-D-Tyr-Lys)

preparation


and targeting integrin receptors in melanoma)

RN 928406-90-0 HCAPLUS


CN Cyclo[L-arginylglycyl-L- $\alpha$ -aspartyl-D-tyrosyl-N6-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-L-lysyl] (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





PAGE 1-B



RN 928406-91-1 HCAPLUS

CN Cyclo[L-arginylglycyl-L- $\alpha$ -aspartyl-D-tyrosyl-N6-[4-[[2-[(1,1-dimethylethoxy)carbonyl]amino]ethyl][2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]-L-lysyl] (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L49 ANSWER 17 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:957931 HCAPLUS Full-text

DOCUMENT NUMBER: 147:202224

TITLE: Metal-based drugs for diagnosis and therapy

AUTHOR(S): Alves, Susana; Vitor, Rute;  
Raposo, Paula D.; Marques, Fernanda; Correia,  
Joao D. G.; Paulo, Antonio;  
Santos, Isabel

CORPORATE SOURCE: Departamento de Quimica, Instituto Tecnologico e Nuclear, Sacavem, 2686-953, Port.

SOURCE: Metal Ions in Biology and Medicine (2006), 9, 3-8  
 CODEN: MIBMCT; ISSN: 1257-2535

PUBLISHER: John Libbey Eurotext

DOCUMENT TYPE: Journal

LANGUAGE: English


OTHER SOURCE(S): CASREACT 147:202224

ED Entered STN: 18 Sep 2006

AB The compound 3,5Me-pz(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub> (L1) is a very effective chelator for the fac-[M(CO)<sub>3</sub>(H<sub>2</sub>O)<sub>3</sub>]<sup>+</sup> (M = Re (1) 99mTc (1a)) moieties, yielding the building blocks fac-[M(CO)<sub>3</sub>(k<sup>3</sup>-L1)]<sup>+</sup> (M = Re (2) 99mTc (2a)). The evaluation of the in vitro and in vivo behavior of 2a showed that this stable building block displays a favorable biol. profile for labeling biomols. with 99mTc, biol. active peptides. Due to its versatility, L1 was integrated through its secondary amine into a peptide with affinity for MC1 receptors (L2), and derivatized with an anthracenyl group at the C(4) position of the pyrazolyl ring (L3). The resulting bifunctional chelators react with 1a yielding the


well defined fac-[<sup>99m</sup>Tc(CO)<sub>3</sub>(k<sup>3</sup>-L)]<sup>+</sup> (L = L<sub>2</sub> (3a), L<sub>3</sub> (4a)) complexes with excellent stability in vitro and in vivo. Complex 3a presents a significant internalization in B16F1 melanoma cells, showing in vivo a significant overall excretion and a reasonable tumor uptake, with a fast clearance from most organs and tissues. For complex 4a, in vitro studies using B16F1 melanoma cells showed significant nuclear internalization and an enhanced radiotoxicity for this compound, most probably due to the presence of the anthracenyl group which is a known DNA intercalator. The results obtained for complexes 3a and 4a indicate that this family of compds. is potentially useful to develop novel specific <sup>99m</sup>Tc radiopharmaceuticals directed for both detection and therapy of melanoma.

- CC 78-7 (Inorganic Chemicals and Reactions)  
Section cross-reference(s): 1
- ST rhenium aminoethylaminoethylpyrazole complex prepn; technetium aminoethylaminoethylpyrazole complex prepn; anthracenylmethyl deriv aminoethylaminoethylpyrazole rhenium technetium complex prepn antitumor activity; peptide sequence pyrazolyl chelator technetium complex prepn antitumor activity
- IT Radiopharmaceuticals  
(preparation of technetium-99m complexes with peptide sequence pyrazolyl chelator and (((aminoethyl)amino)ethyl)pyrazole anthracenylmethyl derivative for development of)
- IT 511513-23-8  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of rhenium and technetium-99m complexes with (((aminoethyl)amino)ethyl)pyrazole and its derivs. with anthracenylmethyl and peptide sequence)
- IT 944389-68-8P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and reaction with technetium-99m aqua carbonyl complex)
- IT 511513-23-8  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(for preparation of rhenium and technetium-99m complexes with (((aminoethyl)amino)ethyl)pyrazole and its derivs. with anthracenylmethyl and peptide sequence)
- RN 511513-23-8 HCAPLUS
- CN 1,2-Ethanediamine, N1-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]- (CA INDEX NAME)




- IT 944389-68-8P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and reaction with technetium-99m aqua carbonyl complex)
- RN 944389-68-8 HCAPLUS
- CN L-Lysinamide, N-acetyl-L-norleucyl-L- $\alpha$ -aspartyl-L-histidyl-L-phenylalanyl-L-arginyl-L-tryptophylglycyl-N-[4-[(2-aminoethyl)[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]amino]-1-oxobutyl]- (CA INDEX NAME)

Absolute stereochemistry.



PAGE 1-B



L49 ANSWER 18 OF 21 HCAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2002:937047 HCAPLUS Full-text  
DOCUMENT NUMBER: 138:330675  
TITLE: Coordination capabilities of pyrazolyl containing  
ligands towards the fac-[Re(CO)3]+ moiety  
AUTHOR(S): Alves, Susana; Paulo, Antonio;  
Correia, Joao D. G.; Domingos, Angela;  
Santos, Isabel  
CORPORATE SOURCE: Departamento de Quimica, ITN, Sacavem, 2686-953, Port.  
SOURCE: Journal of the Chemical Society, Dalton Transactions  
(2002), (24), 4714-4719  
CODEN: JCSDAA; ISSN: 1472-7773  
PUBLISHER: Royal Society of Chemistry  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 138:330675

AB The coordination capabilities of the pyrazolyl containing ligands  $\text{pz}^*(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{pz}^*$ ,  $\text{pz}^*(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{NH}_2$ ,  $\text{pz}^*(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{pz}^*$  and  $\text{pz}^*(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{NH}_2$  ( $\text{pz}^* = 3,5\text{-Me}_2\text{pz}$ ) towards the synthon  $(\text{NEt}_4)_2[\text{ReBr}_3(\text{CO})_3]$  (1) were studied. Depending on the reaction conditions, neutral or cationic  $\text{Re}(\text{I})$  tricarbonyl complexes were isolated:  $[\text{ReBr}(\text{CO})_3(\kappa^2\text{-pz}^*(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{pz}^*)]$  (2),  $[\text{ReBr}(\text{CO})_3(\kappa^2\text{-pz}^*(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{pz}^*)]$  (3)

[Re(CO)<sub>3</sub>(κ<sup>3</sup>-pz\*(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>pz\*)]Br (4), [Re(CO)<sub>3</sub>(κ<sup>2</sup>-pz\*(CH<sub>2</sub>)<sub>2</sub>S(CH<sub>2</sub>)<sub>2</sub>pz\*)MeOH]Br (5), [Re(CO)<sub>3</sub>(κ<sup>3</sup>-pz\*(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub>)]Br (6) and [Re(CO)<sub>3</sub>(κ<sup>3</sup>-pz\*(CH<sub>2</sub>)<sub>2</sub>S(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub>)]Br (7). Complexes 2-7 were characterized by the normal techniques, including x-ray crystallog. anal. in the case of 3, 4, 6 and 7. In these complexes the Re atom adopts a distorted octahedral coordination, being one of the triangular faces defined by the three carbonyl groups and the other three remaining coordination positions by the bidentate and the bromide ligands (3), or by the tridentate and neutral pyrazolyl containing ligands (4, 6, 7). Complexes 2-4, 6 and 7 are static in solution and the <sup>1</sup>H NMR data indicate clearly a κ<sup>2</sup>-coordination mode of the ligand in 2 and 3 and a κ<sup>3</sup>-coordination in 4, 6 and 7, which agrees with the coordination mode found in the solid state. Compound 5 displays a fluxional behavior in solution as shown by variable temperature <sup>1</sup>H NMR studies. No x-ray data exists for this complex but the pattern obtained for the NMR spectrum at 215 K indicates a κ<sup>2</sup>-coordination mode for the pyrazolyl containing ligand.

CC 78-7 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 75

IT 511513-23-8P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and complexation with rhenium carbonyl complex)

IT 511513-23-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and complexation with rhenium carbonyl complex)

RN 511513-23-8 HCAPLUS

CN 1,2-Ethanediamine, N1-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]- (CA INDEX NAME)



REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d ibib ed ab ind 19-21

YOU HAVE REQUESTED DATA FROM FILE 'HCAPLUS, EMBASE, SCISEARCH' - CONTINUE? (Y)/N:y

L49 ANSWER 19 OF 21 EMBASE COPYRIGHT (c) 2009 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 2006052409 EMBASE Full-text

TITLE: Radiopharmaceuticals for targeted radiotherapy.

AUTHOR: Marques, Fernanda (correspondence); Paulo, Antonio  
; Campello, Maria Paula; Lacerda, Sara; Vitor, Rute  
Filipe; Gano, Lurdes; Santos, Isabel

CORPORATE SOURCE: Departamento de Quimica, Instituto Tecnologico e Nuclear,  
EN 10, Apartado 21, 2686-953 Sacavem, Portugal. fmarujo@itn.  
.mces.pt

AUTHOR: Delgado, Rita



CORPORATE SOURCE: Instituto Tecnologia Quimica e Biologica, UNL, Apartado 127, 2781-901 Oeiras, Portugal.

SOURCE: Radiation Protection Dosimetry, (20 Dec 2005) Vol. 116, No. 1-4, pp. 601-604.  
 Refs: 13  
 ISSN: 0144-8420 CODEN: RPDODE

COUNTRY: United Kingdom

DOCUMENT TYPE: Journal; Article

FILE SEGMENT: 016 Cancer  
 023 Nuclear Medicine  
 030 Clinical and Experimental Pharmacology  
 037 Drug Literature Index  
 039 Pharmacy

LANGUAGE: English

SUMMARY LANGUAGE: English

ENTRY DATE: Entered STN: 3 Mar 2006  
 Last Updated on STN: 3 Mar 2006

ED Entered STN: 3 Mar 2006  
 Last Updated on STN: 3 Mar 2006

AB This work intends to find specific radiopharmaceuticals for cancer therapy based on beta ((153)Sm and (166)Ho) or Auger ((99)Tc(m)) emitter radionuclides, using cyclic and acyclic polyamines as bifunctional chelators. These chelators are designed to allow the binding of a tumour seeking biomolecule and/or a DNA intercalator. The cyclic amines, such as 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,8,11-tetraazacyclotetradecane-1,4,8,11-tetraacetic acid and 1,4,7,10-tetraazacyclotridecane-1,4,7,10-tetraacetic acid, were radiolabelled with (153)Sm and (166)Ho. The radiochemical and biological behaviour of the resulting complexes were evaluated in order to assess their potential as building blocks for the attachment of selected biomolecules, with the aim of further applying them for the development of specific therapeutic radiopharmaceuticals. Novel pyrazolyldiamines, bearing a DNA intercalating anthracenyl fragment, were also explored to synthesize radioactive complexes with the fac- (99)Tc(m)(CO) (3) (+) moiety. The identity of these (99)Tc(m) tricarbonyl complexes was confirmed by high-performance liquid chromatography comparison with rhenium congeners fully characterized. By including a DNA intercalator into the chelator framework, we expect to induce more efficient and selective damage to the DNA of cancer cells by the action of the short-range Auger electrons emitted by (99)Tc(m). .COPYRGT. The Author 2005. Published by Oxford University Press. All rights reserved.

CT Medical Descriptors:  
 article  
 beta radiation  
 cancer cell  
 \*cancer radiotherapy  
 clinical trial  
 complex formation  
 DNA damage  
 drug binding  
 drug determination  
 drug selectivity  
 drug structure  
 drug synthesis  
 drug targeting  
 human  
 isotope labeling  
 radiochemistry  
 reversed phase high performance liquid chromatography

CT Drug Descriptors:  
 1,4,7,10 tetraazacyclododecane 1,4,7,10 tetraacetic acid

1,4,7,10 tetraazacyclotridecane 1,4,7,10 tetraacetic acid  
 1,4,8,11 tetraazacyclotetradecane 1,4,8,11 tetraacetic acid  
 anthracene derivative

chelating agent

cyclam derivative

holmium: PR, pharmaceuticals

holmium: PD, pharmacology

holmium 166: PR, pharmaceuticals

holmium 166: PD, pharmacology

intercalating agent

losoxantrone: CT, clinical trial

losoxantrone: AN, drug analysis

losoxantrone: PR, pharmaceuticals

losoxantrone: PD, pharmacology

polyamine derivative

pyrazolyldiamine derivative: PR, pharmaceuticals

pyrazolyldiamine derivative: PD, pharmacology

radioisotope: PR, pharmaceuticals

radioisotope: PD, pharmacology

\*radiopharmaceutical agent: CT, clinical trial

\*radiopharmaceutical agent: AN, drug analysis

\*radiopharmaceutical agent: PR, pharmaceuticals

\*radiopharmaceutical agent: PD, pharmacology

rhenium complex

samarium 153: PR, pharmaceuticals

samarium 153: PD, pharmacology

technetium 99m: PR, pharmaceuticals

technetium 99m: PD, pharmacology

unclassified drug

RN (1,4,7,10 tetraazacyclododecane 1,4,7,10 tetraacetic acid) 60239-18-1;  
 (holmium) 7440-60-0; (losoxantrone) 88303-60-0; (samarium 153) 15766-00-4;  
 (technetium 99m) 14133-76-7

L49 ANSWER 20 OF 21 SCISEARCH COPYRIGHT (c) 2009 The Thomson Corporation on  
 STN

ACCESSION NUMBER: 2009:366412 SCISEARCH Full-text

THE GENUINE ARTICLE: 417GQ

TITLE: Influence of the ligand donor atoms on the in vitro  
 stability of rhenium(I) and technetium (I)-99m complexes  
 with pyrazole-containing chelators:  
 Experimental and DFT studies

AUTHOR: Santos, Isabel (Reprint)

CORPORATE SOURCE: ITN, Unidade Ciencias Quim & Radiofarmaceut, Estr Nacl 10,  
 P-2686953 Sacavem Codex, Portugal (Reprint)  
 E-mail: isantos@itn.pt

AUTHOR: Santos, Isabel (Reprint)

CORPORATE SOURCE: ITN, Unidade Ciencias Quim & Radiofarmaceut, P-2686953  
 Sacavem Codex, Portugal  
 E-mail: isantos@itn.pt

AUTHOR: Moura, Carolina; Fernandes, Celia; Gano, Lurdes;  
Paulo, Antonio; Santos, Isabel C.;  
 Calhorda, Maria Jose

CORPORATE SOURCE: Univ Lisbon, Dept Quim & Bioquim, CQB, Fac Ciencias,  
 P-1749016 Lisbon, Portugal

COUNTRY OF AUTHOR: Portugal

SOURCE: JOURNAL OF ORGANOMETALLIC CHEMISTRY, (15 MAR 2009) Vol.  
 694, No. 6, pp. 950-958.  
 ISSN: 0022-328X.

PUBLISHER: ELSEVIER SCIENCE SA, PO BOX 564, 1001 LAUSANNE,  
 SWITZERLAND.

DOCUMENT TYPE: Article; Journal  
 LANGUAGE: English  
 REFERENCE COUNT: 63  
 ENTRY DATE: Entered STN: 26 Mar 2009

Last Updated on STN: 7 May 2009

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

ED Entered STN: 26 Mar 2009

Last Updated on STN: 7 May 2009

AB The new pyrazole-containing ligand 3,5-Me(2)pz(CH<sub>2</sub>)(2)S(CH<sub>2</sub>)(2)COOH ((LH)-H-1) was synthesized and used to prepare the complexes fac-[M(kappa(3)-L-1)(CO)(3)] (M = Re (1), (99mTc)(1a)), which were obtained in high yield albeit with a low specific activity in the case of Tc-99m. The X-ray diffraction analysis of 1 confirmed that L-1 coordinates to the metal as monoanionic and through a (N,S,O) donor atom set. Challenge experiments of 1a against cysteine and histidine showed that this complex suffers considerable transchelation in vitro. This contrasts with the behavior exhibited by the related complex fac-[Tc-99m(kappa(3)-L-2)(CO)(3)] (2a) (L-2 = 3,5-Me(2)pz-(CH<sub>2</sub>)(2)NH-CH<sub>2</sub>-COO), anchored by a (N2O)-tridentate ligand. Biodistribution studies of 1a and 2a in mice indicated that both compounds have a relatively similar biological profile. Nevertheless, the fastest blood clearance and minor hepatic retention found for 2a has shown that this complex is more adequate to be further explored in radiopharmaceutical sciences. DFT calculations (ADF program) were performed for these neutral complexes and related cationic M(I) (M = Re, Tc) tricarbonyl complexes anchored by pyrazole-containing ligands, in order to have a better understanding of the influence of the donor atom set (N,N,O vs. N,O,S; N,N,N vs. N,N,S vs. N,S,S) on their in vitro stability. The differences of the calculated binding energies are not significant, suggesting that the in vitro behavior of these Re(I)/Tc(I) tricarbonyl complexes is not determined by thermodynamic factors. (C) 2008 Elsevier B. V. All rights reserved.

CC CHEMISTRY, INORGANIC & NUCLEAR; CHEMISTRY, ORGANIC

ST Author Keywords: Rhenium; Technetium; Carbonyl; Pyrazolyl-containing ligands; DFT calculations

STP KeyWords Plus (R): TRANSITION-STATE METHOD; REGULAR 2-COMPONENT HAMILTONIANS; DENSITY-FUNCTIONAL THEORY; TRICARBONYL COMPLEXES; SCREENING MODEL; VIVO EVALUATION; RADIOPHARMACEUTICALS; APPROXIMATION; BIOMOLECULES; DERIVATIVES

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

L49 ANSWER 21 OF 21 SCISEARCH COPYRIGHT (c) 2009 The Thomson Corporation on STN

ACCESSION NUMBER: 1999:482083 SCISEARCH Full-text

THE GENUINE ARTICLE: 209HU

TITLE: Derivative chemistry of [UC12{B(pz)(4)}(2)]: stability of complexes containing the fragments [U{B(pz)(4)}(2)] and [U{HB(pz)(3)}(2)]

AUTHOR: Santos I (Reprint)

CORPORATE SOURCE: ITN, Dept Quim, P-2686 Sacavem, Portugal (Reprint)

AUTHOR: Campello M P C; Domingos A; Galvao A; de Matos A P

CORPORATE SOURCE: Inst Super Tecn, Dept Engrn Quim, P-1096 Lisbon, Portugal

COUNTRY OF AUTHOR: Portugal

SOURCE: JOURNAL OF ORGANOMETALLIC CHEMISTRY, (5 MAY 1999) Vol. 579, No. 1-2, pp. 5-17.  
 ISSN: 0022-328X.

PUBLISHER: ELSEVIER SCIENCE SA, PO BOX 564, 1001 LAUSANNE, SWITZERLAND.

DOCUMENT TYPE: Article; Journal

LANGUAGE: English

REFERENCE COUNT: 42

ENTRY DATE: Entered STN: 1999  
 Last Updated on STN: 1999  
 \*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

ED Entered STN: 1999  
 Last Updated on STN: 1999

AB Uranium tetrachloride reacts with two equivalents of K[B(pz)(4)] in THF affording [UCl<sub>2</sub>{B(pz)(4)}(2)] (1) in 75% yield. Complex 1 is monomeric and crystallizes in the monoclinic space group C2/c with cell parameters a = 13.700(6), b = 12.759(2), c = 17.513(8) Angstrom, beta = 101.37(2)degrees, V = 3001(2) Angstrom(3), Z = 4. Derivatives [UCl(OR){B(pz)(4)}(2)] (R=C<sub>2</sub>H<sub>5</sub> (2), Bu-t (3), C<sub>6</sub>H<sub>4</sub>-o-OMe (4) and C<sub>6</sub>H<sub>2</sub>-2,4,6-Me-3 (5)), [U((OBu)-Bu-t)(2){B(pz)(4)}(2)] (6), [U((SPr)-Pr-i)(2){B(pz)(4)}(2)] (7) and [UCl(Me){B(pz)(4)}(2)] (8) were obtained by reacting 1 with sodium alkoxides, with (NaSPr)-Pr-i or with LiMe. X-ray crystallographic analysis of 5 and 7 shows that uranium is eight-coordinate by the two eta(3)-[B(pz)(4)] ligands and by two monodentate coligands (5. crystallizes in the monoclinic space group C2/c with cell parameters a = 30.575(3). b = 9.929(1), c = 24.884(3) Angstrom, beta = 90.59(1)degrees, V = 7554(1) Angstrom(3), Z = 8; 7 crystallizes in the monoclinic space group C2/c with cell parameters a = 24.286(7), b = 9.471(2), c = 16.076(3) Angstrom, beta = 96.44(3)degrees, V = 3674(2) Angstrom(3), Z = 4). Extended Huckel molecular orbital (EHMO) calculations were used to get a better insight into the electronic properties of the ligand [B(pz)(4)] and to get some explanation on the relative stability of complexes containing the fragments '[U(B(pz)(4)}(2)] and '[U(HB(pz)(3)}(2)]'. (C) 1999 Elsevier Science S.A. All rights reserved.

CC CHEMISTRY, INORGANIC & NUCLEAR; CHEMISTRY, ORGANIC

ST Author Keywords: uranium; poly(pyrazolyl)borates; EHMO calculations; stability

STP KeyWords Plus (R): ION SIZE DISCRIMINATION; X-RAY CRYSTAL; POLY(PYRAZOLYL)BORATE LIGANDS; MOLECULAR-STRUCTURES; CHELATE COMPLEXES; LANTHANIDE IONS; INTRALIGAND; CONTACT; MODE  
 \*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

=> file stnguide

FILE 'STNGUIDE' ENTERED AT 13:45:22 ON 25 JUN 2009  
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
 COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Jun 19, 2009 (20090619/UP).

=> d his ful

(FILE 'HOME' ENTERED AT 12:43:21 ON 25 JUN 2009)

FILE 'STNGUIDE' ENTERED AT 12:43:24 ON 25 JUN 2009

FILE 'ZCAPLUS' ENTERED AT 12:44:29 ON 25 JUN 2009  
E US2005-551292/APPS

L1 FILE 'HCAPLUS' ENTERED AT 12:44:45 ON 25 JUN 2009  
1 SEA SPE=ON ABB=ON PLU=ON US2005-551292/APPS  
D SCAN

FILE 'STNGUIDE' ENTERED AT 12:45:05 ON 25 JUN 2009

L2 FILE 'WPIX' ENTERED AT 12:45:26 ON 25 JUN 2009  
1 SEA SPE=ON ABB=ON PLU=ON US2005-551292/APPS  
D IALL CODE

FILE 'STNGUIDE' ENTERED AT 12:46:49 ON 25 JUN 2009

FILE 'REGISTRY' ENTERED AT 12:47:09 ON 25 JUN 2009

L3 FILE 'HCAPLUS' ENTERED AT 12:47:12 ON 25 JUN 2009  
TRA PLU=ON L1 1- RN : 29 TERMS

L4 FILE 'REGISTRY' ENTERED AT 12:47:14 ON 25 JUN 2009  
29 SEA SPE=ON ABB=ON PLU=ON L3  
D SCAN

FILE 'STNGUIDE' ENTERED AT 12:47:38 ON 25 JUN 2009

L5 FILE 'LREGISTRY' ENTERED AT 12:49:19 ON 25 JUN 2009  
STR

L6 FILE 'REGISTRY' ENTERED AT 12:54:28 ON 25 JUN 2009  
5 SEA SSS SAM L5

FILE 'STNGUIDE' ENTERED AT 12:54:53 ON 25 JUN 2009  
D QUE STAT

L7 FILE 'REGISTRY' ENTERED AT 12:59:40 ON 25 JUN 2009  
827 SEA SSS FUL L5  
SAVE TEMP L7 SCH292PSET1/A

L8 10 SEA SPE=ON ABB=ON PLU=ON L4 AND L7

L9 19 SEA SPE=ON ABB=ON PLU=ON L4 NOT L8  
D SCAN

FILE 'STNGUIDE' ENTERED AT 13:01:33 ON 25 JUN 2009

L10 FILE 'LREGISTRY' ENTERED AT 13:02:03 ON 25 JUN 2009  
STR L5

L11 FILE 'REGISTRY' ENTERED AT 13:03:00 ON 25 JUN 2009  
2 SEA SUB=L7 SSS SAM L10  
D SCAN

L12 90 SEA SUB=L7 SSS FUL L10  
SAVE TEMP L12 SCH292RSET1/A

L13 0 SEA SPE=ON ABB=ON PLU=ON L8 NOT L12

FILE 'STNGUIDE' ENTERED AT 13:05:03 ON 25 JUN 2009

FILE 'REGISTRY' ENTERED AT 13:06:34 ON 25 JUN 2009

FILE 'ZCAPLUS' ENTERED AT 13:07:06 ON 25 JUN 2009

```

L*** DEL      QUE SANTOS, I?/AU
L*** DEL      QUE REGO, I?/AU
L14          QUE SPE=ON  ABB=ON  PLU=ON  SANTOS, I?/AU,AUTH
L15          QUE SPE=ON  ABB=ON  PLU=ON  REGO, I?/AU,AUTH
L16          QUE SPE=ON  ABB=ON  PLU=ON  CORREIA, J?/AU,AUTH
L17          QUE SPE=ON  ABB=ON  PLU=ON  PAULO, A?/AU,AUTH
L18          QUE SPE=ON  ABB=ON  PLU=ON  ALVES, S?/AU,AUTH
L19          QUE SPE=ON  ABB=ON  PLU=ON  VITOR, R?/AU,AUTH
L20          QUE SPE=ON  ABB=ON  PLU=ON  MALLINCKRODT/CS,SO,PA
L21          QUE SPE=ON  ABB=ON  PLU=ON  AY<2004 OR PY<2004 OR PRY<2004 OR
MY<2004 OR REVIEW/DT
L22          QUE SPE=ON  ABB=ON  PLU=ON  BIOMOLECUL? OR (BIO(1W)MOLECUL?)
OR (BIOLOGIC?(3A)MOLECUL?)
L23          QUE SPE=ON  ABB=ON  PLU=ON  CHELAT?
L24          QUE SPE=ON  ABB=ON  PLU=ON  "CHELATING AGENTS"+PFT,OLD,NEW,NT/C
T

```

FILE 'HCAPLUS' ENTERED AT 13:21:41 ON 25 JUN 2009

```

L25          46 SEA SPE=ON  ABB=ON  PLU=ON  L12
L26          19 SEA SPE=ON  ABB=ON  PLU=ON  L25 AND (L22 OR L23 OR L24)
L27          46 SEA SPE=ON  ABB=ON  PLU=ON  (L25 OR L26)
L28          18 SEA SPE=ON  ABB=ON  PLU=ON  L27 AND (L14 OR L15 OR L16 OR L17
OR L18 OR L19 OR L20)
L29          28 SEA SPE=ON  ABB=ON  PLU=ON  L27 NOT L28
L30          23 SEA SPE=ON  ABB=ON  PLU=ON  L29 AND L21
L31          28 SEA SPE=ON  ABB=ON  PLU=ON  (L29 OR L30)

```

FILE 'WPIX' ENTERED AT 13:24:10 ON 25 JUN 2009

```

D QUE L12
L32          2 SEA SSS SAM L10
D TRI 1-2
L33          10 SEA SSS FUL L10
SAVE TEMP L33 SCH292WPIS/A
SELECT L33 1- SDCN
L34          3 SEA SPE=ON  ABB=ON  PLU=ON  (RABNX7/DCN OR RAFVJB/DCN OR
RAFVJC/DCN OR RAFVJD/DCN OR RAFVJE/DCN OR RAFVJF/DCN OR
RAFVJG/DCN OR RAFVJ8/DCN OR RAFVJ9/DCN OR RAMT8E/DCN) OR
L33/DCR
L35          2 SEA SPE=ON  ABB=ON  PLU=ON  L34 AND (L22 OR L23)
L36          3 SEA SPE=ON  ABB=ON  PLU=ON  (L34 OR L35)
L37          1 SEA SPE=ON  ABB=ON  PLU=ON  L36 AND (L14 OR L15 OR L16 OR L17
OR L18 OR L19 OR L20)
L38          2 SEA SPE=ON  ABB=ON  PLU=ON  L36 NOT L37
D TRI HITSTR 1-2

```

FILE 'STNGUIDE' ENTERED AT 13:27:39 ON 25 JUN 2009

FILE 'MEDLINE' ENTERED AT 13:28:54 ON 25 JUN 2009

```

L39          0 SEA SPE=ON  ABB=ON  PLU=ON  L12

```

FILE 'REGISTRY' ENTERED AT 13:29:10 ON 25 JUN 2009

```

L40          10 SEA SPE=ON  ABB=ON  PLU=ON  L4 AND L12

```

FILE 'MEDLINE' ENTERED AT 13:29:34 ON 25 JUN 2009

10/551,292

FILE 'REGISTRY' ENTERED AT 13:29:40 ON 25 JUN 2009  
SET SMARTSELECT ON  
SET SMARTSELECT OFF

FILE 'MEDLINE' ENTERED AT 13:29:40 ON 25 JUN 2009

FILE 'EMBASE' ENTERED AT 13:29:56 ON 25 JUN 2009  
L41 0 SEA SPE=ON ABB=ON PLU=ON L12

FILE 'STNGUIDE' ENTERED AT 13:30:05 ON 25 JUN 2009

FILE 'BIOSIS, CABA, AGRICOLA, BIOTECHNO, DRUGU, VETU' ENTERED AT 13:30:22  
ON 25 JUN 2009  
L42 0 SEA SPE=ON ABB=ON PLU=ON L12

FILE 'STNGUIDE' ENTERED AT 13:30:39 ON 25 JUN 2009

FILE 'MEDLINE, BIOSIS, EMBASE, PASCAL, JAPIO, CABA, CEABA-VTB, LIFESCI,  
BIOENG, BIOTECHNO, BIOTECHDS, DRUGU, DRUGB, VETU, VETB, SCISEARCH,  
CONFSCI, DISSABS, RDISCLOSURE' ENTERED AT 13:31:47 ON 25 JUN 2009  
L43 769 SEA SPE=ON ABB=ON PLU=ON ?PYRAZOL?/IT, TI, CC, CT, ST, STP AND  
L23/IT, TI, CC, CT, ST, STP  
L44 20 SEA SPE=ON ABB=ON PLU=ON L43 AND (L14 OR L15 OR L16 OR L17  
OR L18 OR L19 OR L20)

FILE 'STNGUIDE' ENTERED AT 13:33:10 ON 25 JUN 2009  
D QUE STAT L7  
D QUE STAT L12  
D QUE NOS L31  
D QUE STAT L33  
D QUE NOS L38  
D QUE NOS L39  
D QUE NOS L41  
D QUE NOS L42

FILE 'HCAPLUS, WPIX' ENTERED AT 13:37:35 ON 25 JUN 2009  
L45 28 DUP REM L31 L38 L39 L41 L42 (2 DUPLICATES REMOVED)  
ANSWERS '1-27' FROM FILE HCAPLUS  
ANSWER '28' FROM FILE WPIX  
SAVE TEMP L45 SCH292MAINP/A

FILE 'STNGUIDE' ENTERED AT 13:37:48 ON 25 JUN 2009

FILE 'HCAPLUS, WPIX' ENTERED AT 13:38:10 ON 25 JUN 2009  
D QUE L21  
L\*\*\* DEL 28 S L29-L30  
L\*\*\* DEL 2 S L36 NOT L37  
L46 23 SEA SPE=ON ABB=ON PLU=ON L45 AND L21  
L47 23 DUP REM L46 (0 DUPLICATES REMOVED)  
ANSWERS '1-22' FROM FILE HCAPLUS  
ANSWER '23' FROM FILE WPIX

FILE 'STNGUIDE' ENTERED AT 13:39:14 ON 25 JUN 2009

FILE 'HCAPLUS, WPIX' ENTERED AT 13:39:20 ON 25 JUN 2009  
D IBIB ED ABS HITIND HITSTR 1-22

FILE 'STNGUIDE' ENTERED AT 13:39:37 ON 25 JUN 2009

10/551,292

FILE 'HCAPLUS, WPIX' ENTERED AT 13:41:17 ON 25 JUN 2009  
D IALL ABEQ TECH ABEX FRAGHITSTR 23

FILE 'STNGUIDE' ENTERED AT 13:41:18 ON 25 JUN 2009

FILE 'HCAPLUS, WPIX' ENTERED AT 13:41:34 ON 25 JUN 2009

L\*\*\* DEL 28 S L29-L30  
L\*\*\* DEL 22 S L45 AND L21  
L\*\*\* DEL 2 S L36 NOT L37  
L\*\*\* DEL 1 S L45 AND L21  
L48 5 SEA SPE=ON ABB=ON PLU=ON L45 NOT L47

FILE 'STNGUIDE' ENTERED AT 13:41:51 ON 25 JUN 2009

FILE 'HCAPLUS' ENTERED AT 13:41:55 ON 25 JUN 2009  
D IBIB ED ABS HITIND HITSTR 1-5

FILE 'STNGUIDE' ENTERED AT 13:42:02 ON 25 JUN 2009

D QUE NOS L28  
D QUE NOS L37  
D QUE NOS L39  
D QUE NOS L41  
D QUE NOS L42  
D QUE L44

FILE 'HCAPLUS, WPIX, MEDLINE, BIOSIS, EMBASE, PASCAL, BIOENG, SCISEARCH'  
ENTERED AT 13:43:20 ON 25 JUN 2009

L49 21 DUP REM L28 L37 L39 L41 L42 L44 (18 DUPLICATES REMOVED)  
ANSWERS '1-18' FROM FILE HCAPLUS  
ANSWER '19' FROM FILE EMBASE  
ANSWERS '20-21' FROM FILE SCISEARCH  
SAVE TEMP L49 SCH292INV/A

FILE 'STNGUIDE' ENTERED AT 13:43:36 ON 25 JUN 2009

FILE 'HCAPLUS, EMBASE, SCISEARCH' ENTERED AT 13:43:55 ON 25 JUN 2009  
D IBIB ED ABS HITIND HITSTR 1-18

FILE 'STNGUIDE' ENTERED AT 13:44:05 ON 25 JUN 2009

FILE 'HCAPLUS, EMBASE, SCISEARCH' ENTERED AT 13:45:06 ON 25 JUN 2009  
D IBIB ED AB IND 19-21

FILE 'STNGUIDE' ENTERED AT 13:45:07 ON 25 JUN 2009

FILE 'STNGUIDE' ENTERED AT 13:45:22 ON 25 JUN 2009

FILE HOME

FILE STNGUIDE  
FILE CONTAINS CURRENT INFORMATION.  
LAST RELOADED: Jun 19, 2009 (20090619/UP).

FILE ZCAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.



10/551,292

The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS is strictly prohibited.

FILE COVERS 1907 - 25 Jun 2009 VOL 150 ISS 26  
FILE LAST UPDATED: 24 Jun 2009 (20090624/ED)  
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Apr 2009  
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Apr 2009

ZCAplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE HCAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 25 Jun 2009 VOL 150 ISS 26  
FILE LAST UPDATED: 24 Jun 2009 (20090624/ED)  
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Apr 2009  
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Apr 2009

HCAplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE WPIX

FILE LAST UPDATED: 19 JUN 2009 <20090619/UP>  
MOST RECENT UPDATE: 200939 <200939/DW>  
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE  
>>> Now containing more than 1.4 million chemical structures in DCR <<<

>>> IPC, ECLA and US National Classifications have been updated with reclassifications to March 15th, 2009.

F-Term and FI-Term original classifications are current and reclassification will commence in June.

No update date (UP) has been created for the reclassified documents, but they can be identified by

10/551,292

specific update codes (see HELP CLA for details)<<<

FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,  
PLEASE VISIT:

[http://www.stn-international.com/stn\\_guide.html](http://www.stn-international.com/stn_guide.html)

FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE

<http://scientific.thomsonreuters.com/support/patents/coverage/latestupdate>

EXPLORE DERWENT WORLD PATENTS INDEX IN STN ANAVIST, VERSION 2.0:

[http://www.stn-international.com/DWPIAnaVist2\\_0608.html](http://www.stn-international.com/DWPIAnaVist2_0608.html)

>>> HELP for European Patent Classifications see HELP ECLA, HELP ICO <<<

#### FILE REGISTRY

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 23 JUN 2009 HIGHEST RN 1159631-40-9

DICTIONARY FILE UPDATES: 23 JUN 2009 HIGHEST RN 1159631-40-9

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 9, 2009.

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and  
predicted properties as well as tags indicating availability of  
experimental property data in the original document. For information  
on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

#### FILE LREGISTRY

LREGISTRY IS A STATIC LEARNING FILE

CAS INFORMATION USE POLICIES, ENTER HELP USAGETERMS FOR DETAILS.

#### FILE MEDLINE

FILE LAST UPDATED: 24 Jun 2009 (20090624/UP). FILE COVERS 1949 TO DATE.

MEDLINE and LMEDLINE have been updated with the 2009 Medical Subject  
Headings (MeSH) vocabulary and tree numbers from the U.S. National Library  
of Medicine (NLM). Additional information is available at

[http://www.nlm.nih.gov/pubs/techbull/nd08/nd08\\_medline\\_data\\_changes\\_2009](http://www.nlm.nih.gov/pubs/techbull/nd08/nd08_medline_data_changes_2009).

On February 21, 2009, MEDLINE was reloaded. See HELP RLOAD for details.

This file contains CAS Registry Numbers for easy and accurate  
substance identification.

See HELP RANGE before carrying out any RANGE search.

#### FILE EMBASE

FILE COVERS 1974 TO 25 Jun 2009 (20090625/ED)

10/551,292

EMBASE was reloaded on March 30, 2008.

EMBASE is now updated daily. SDI frequency remains weekly (default) and biweekly.

This file contains CAS Registry Numbers for easy and accurate substance identification.

Beginning January 2008, Elsevier will no longer provide EMTREE codes as part of the EMTREE thesaurus in EMBASE. Please update your current-awareness alerts (SDIs) if they contain EMTREE codes.

For further assistance, please contact your local helpdesk.

FILE BIOSIS

FILE COVERS 1926 TO DATE.

CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNs) PRESENT  
FROM JANUARY 1926 TO DATE.

RECORDS LAST ADDED: 24 June 2009 (20090624/ED)

BIOSIS has been augmented with 1.8 million archival records from 1926 through 1968. These records have been re-indexed to match current BIOSIS indexing.

FILE CABA

FILE COVERS 1973 TO 4 Jun 2009 (20090604/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

The CABA file was reloaded 7 December 2003. Enter HELP RLOAD for details.

FILE AGRICOLA

FILE COVERS 1970 TO 10 Jun 2009 (20090610/ED)

Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted material. All rights reserved. (2009)

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE BIOTECHNO

FILE LAST UPDATED: 7 JAN 2004 <20040107/UP>

FILE COVERS 1980 TO 2003.

THIS FILE IS A STATIC FILE WITH NO UPDATES

>>> SIMULTANEOUS LEFT AND RIGHT TRUNCATION AVAILABLE IN  
/CT AND BASIC INDEX <<<

FILE DRUGU

FILE LAST UPDATED: 24 JUN 2009 <20090624/UP>

>>> DERWENT DRUG FILE (SUBSCRIBER) <<<

>>> FILE COVERS 1983 TO DATE <<<

>>> THESAURUS AVAILABLE IN /CT <<<

FILE VETU  
FILE LAST UPDATED: 2 JAN 2002 <20020102/UP>  
FILE COVERS 1983-2001

FILE PASCAL  
FILE LAST UPDATED: 22 JUN 2009 <20090622/UP>  
FILE COVERS 1977 TO DATE.

>>> SIMULTANEOUS LEFT AND RIGHT TRUNCATION IS AVAILABLE  
IN THE BASIC INDEX (/BI) FIELD <<<

FILE JAPIO  
FILE LAST UPDATED: 8 JUN 2009 <20090608/UP>  
MOST RECENT PUBLICATION DATE: 26 FEB 2009 <20090226/PD>

>>> GRAPHIC IMAGES AVAILABLE <<<

FILE CEABA-VTB  
FILE LAST UPDATED: 25 JUN 2009 <20090625/UP>  
FILE COVERS 1966 TO DATE

>>> DECHEMA, the producer of CEABA-VTB is using a new classification  
scheme.  
The new classification schemes are available as a PDF file  
and may be downloaded free-of-charge from:  
<http://www.stn-international.com/cc-de.html>  
and  
<http://www.stn-international.com/cc-en.html><<<

FILE LIFESCI  
FILE COVERS 1978 TO 1 May 2009 (20090501/ED)

FILE BIOENG  
FILE LAST UPDATED: 3 JUN 2009 <20090603/UP>  
FILE COVERS 1982 TO DATE

>>> SIMULTANEOUS LEFT AND RIGHT TRUNCATION AVAILABLE IN  
THE BASIC INDEX <<<

FILE BIOTECHDS  
FILE LAST UPDATED: 19 JUN 2009 <20090619/UP>  
FILE COVERS 1982 TO DATE

>>> USE OF THIS FILE IS LIMITED TO BIOTECH SUBSCRIBERS <<<

FILE DRUGB  
>>> FILE COVERS 1964 TO 1982 - CLOSED FILE <<<

FILE VETB  
FILE LAST UPDATED: 25 SEP 94 <940925/UP>  
FILE COVERS 1968-1982

FILE SCISEARCH

10/551,292

FILE COVERS 1974 TO 18 Jun 2009 (20090618/ED)

SCISEARCH has been reloaded, see HELP RLOAD for details.

FILE CONFSCI

FILE COVERS 1973 TO 30 Mar 2009 (20090330/ED)

CSA has resumed updates, see NEWS FILE

FILE DISSABS

FILE COVERS 1861 TO 28 MAY 2009 (20090528/ED)

Only fair use as provided by the United States copyright law is permitted. PROQUEST INFORMATION AND LEARNING COMPANY MAKES NO WARRANTY REGARDING THE ACCURACY, COMPLETENESS OR TIMELINESS OF THE LICENSED MATERIALS OR ANY WARRANTY, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND SHALL NOT BE LIABLE FOR DAMAGES OF ANY KIND OR LOST PROFITS OR OTHER CLAIMS RELATED TO THE LICENSED MATERIALS OR THEIR USE.

FILE RDISCLOSURE

FILE LAST UPDATED: 15 JUN 2009 <20090615/UP>

FILE COVERS 1960 TO DATE

>>> SIMULTANEOUS LEFT AND RIGHT TRUNCATION IS AVAILABLE IN THE  
BASIC INDEX (/BI) AND TITLE (/TI) FIELDS <<<

>>> IMAGES ARE AVAILABLE ONLINE AND FOR EMAIL-PRINTS <<<

=>